

A Z -set unknotting theorem for Nöbeling spaces

Michael Levin

Abstract

We prove a Z -set unknotting theorem for Nöbeling spaces. This generalizes a result obtained by S. Ageev for a restricted class of Z -sets. The theorem is proved for a certain model of Nöbeling spaces.

Keywords: Z -set unknotting theorem, Nöbeling spaces

Math. Subj. Class.: 55M10, 54F45.

1 Introduction

All spaces are assumed to be separable metrizable. A manifold means a manifold with (possibly empty) boundary and a triangulated space means a locally finite simplicial complex which we identify with the underlying space. For a triangulated space we consider only triangulations compatible with the PL-structure of the space. All triangulated manifolds are assumed to be combinatorial.

A complete n -dimensional metric space X is said to be an n -dimensional Nöbeling space if the following properties are satisfied:

- (i) X is an absolute extensor in dimension n , that is every map $f : A \longrightarrow X$ from a closed subset A of a space Y of $\dim \leq n$ extends over Y ;
- (ii) every map $f : Y \longrightarrow X$ from a complete metric space Y of $\dim \leq n$ can be arbitrarily closely approximated by a closed embedding, that is for every open cover $\mathcal{U}$ of X there is a closed embedding $g : Y \longrightarrow X$ which is $\mathcal{U}$ -close to f ($\mathcal{U}$ -close means that for every $y \in Y$ there is an element of $\mathcal{U}$ that contains both $f(y)$ and $g(y)$).

Examples of Nöbeling spaces can be constructed as follows.

By a rational map between triangulated spaces we mean a PL-map that sends points with rational barycentric coordinates to points with rational barycentric coordinates. Two triangulations of a space are said to be rationally equivalent if the identity map is a rational map with respect to these triangulations (it is easy to see that if a PL-homeomorphism is rational in one direction then it is rational in the opposite direction as well). Let M be a triangulated space. Every triangulation of M which is rationally equivalent to the given triangulation of M is said to be a rational triangulation and the class of all rational

triangulations is said to be the rational structure of M . Denote by $M(k)$ the subspace of M which is the complement of the union of all the triangulated spaces of $\dim \leq k$ which are rationally embedded in M .

The space $M(k)$ admits the following interpretation. Fix a rational triangulation of M and embed M in the Hilbert space H by an embedding which sends the vertices to points with rational coordinates and which is linear on every simplex of the triangulation. A k -dimensional plane in H is said to be rational if it is spanned by $k + 1$ points with rational coordinates. Denote by K the union of all rational k -dimensional planes. Then $M(k) = M \setminus K$. Indeed, for every simplex Δ of M and every rational k -dimensional plane L , $\Delta \cap L$ admits a triangulation for which it is rationally embedded in M and therefore $M(k) \subset M \setminus K$. Now let $e : \Delta' \rightarrow M$ be a rational embedding of a simplex Δ' of $\dim \leq k$. Then there is a (not necessarily rational) triangulation $\mathcal{T}$ of Δ' such that e is linear on every simplex of $\mathcal{T}$. Since every $\Delta'' \in \mathcal{T}$ with $\dim \Delta'' = \dim \Delta'$ has a dense subset of points with rational barycentric coordinates with respect to Δ' we conclude that $e(\Delta'')$ is contained in a k -dimensional rational plane in H . Thus $e(\Delta') \subset K$ and hence $M(k) = M \setminus K$.

Let us state the following important fact leaving its proof to the reader.

Theorem 1.1 *Let M be a triangulated m -dimensional manifold, let $k \geq 0$ be an integer and let $n = m - k - 1$. If M is $(n - 1)$ -connected and $m \geq 2n + 1$ then $M(k)$ is an n -dimensional Nöbeling space.*

A space $M(k)$ satisfying the assumptions of Theorem 1.2 will be called a Nöbeling space modeled on a triangulated manifold.

A subset A of a space X is called a Z -set if A is closed in X and the identity map of X can be arbitrarily closely approximated by a map $f : X \rightarrow X$ with $f(X) \cap A = \emptyset$. Note that if X is an n -dimensional Nöbeling space modeled on a manifold M and $A \subset X$ is a Z -set in X then $X \setminus A$ is also an n -dimensional Nöbeling space modeled on the manifold $N = M \setminus \text{the closure of } A \text{ in } M$ (the rational structure of N is defined such that the inclusion is a rational map).

The main result of this paper is

Theorem 1.2 *Let X_1 and X_2 be n -dimensional Nöbeling spaces and let A_1 and A_2 be Z -sets in X_1 and X_2 respectively such that $X_1 \setminus A_1$ and $X_2 \setminus A_2$ are homeomorphic to n -dimensional Nöbeling spaces modeled on triangulated manifolds. If A_1 and A_2 are homeomorphic then any homeomorphism between A_1 and A_2 can be extended to a homeomorphism between X_1 and X_2 .*

Theorem 1.2 was proved by Ageev [1] under some additional restrictions on the sets A_1 and A_2 . The proof of Theorem 1.2 is self-contained and relies only on well-known facts of PL-topology [4, 7], Nöbeling spaces [3] and elementary properties of partitions presented in the very beginning of [2]. Some ideas of the proof of Theorem 1.2 came from [5].

The results of this paper along with some additional arguments can be applied to validate the characterization theorem for Nöbeling saying that any two Nöbeling spaces

of the same dimension are homeomorphic. A proof of the characterization theorem will be presented in a forthcoming paper. Note that the characterization theorem implies that the assumptions on the complements $X_1 \setminus A_1$ and $X_2 \setminus A_2$ in Theorem 1.2 are automatically satisfied and therefore can be dropped.

Let us finally mention that both a Z -set unknotting theorem and the characterization theorem for Nöbeling spaces were also announced by A. Nagorko at International Conference on Geometric Topology, July 2005, Bedlewo, Poland.

2 Preliminaries

2.1 General

Let M be a triangulated manifold. The geometric interior $\text{Int}M$ of M is the set of points having a neighborhood PL-homeomorphic to an Euclidean space of dimension $= \dim M$. The set $\partial M = M \setminus \text{Int}M$ is the geometric boundary of M .

A triangulated space (manifold) PL-embedded in M is said to be a PL-subspace (PL-submanifold) of M . An open subset of a triangulated space is always considered with the induced PL-structure for which the inclusion is a PL-map. Note that an open subset of a PL-submanifold of M is also a PL-submanifold of M .

A subset P of M is said to be a PL-subcomplex of M if there is a triangulation of M for which P is a subcomplex. A subset R of M is said to be a PL-presented subset of M if there are closed subsets $R_1 \subset \dots \subset R_n$ of M such that $R_n = R$, R_1 is a PL-subcomplex of M and $R_{i+1} \setminus R_i$ is a PL-subcomplex of $M \setminus R_i$, $i = 1, \dots, n-1$.

A collection $\mathcal{P}$ of subsets of M is said to be a decomposition of M if $\mathcal{P}$ is a locally finite cover of M and the elements of $\mathcal{P}$ are PL-subcomplexes of M . By a finite intersection of a decomposition $\mathcal{P}$ we mean an intersection of finitely many elements of $\mathcal{P}$ (the elements of $\mathcal{P}$ are also considered as finite intersections). Note that since $\mathcal{P}$ is locally finite, any non-empty intersection of elements of $\mathcal{P}$ must be a finite intersection of $\mathcal{P}$. It is clear that for a PL-submanifold N of M , the restriction $\mathcal{P}|N = \{P \cap N : P \in \mathcal{P}\}$ of $\mathcal{P}$ to N is a decomposition of N .

A decomposition $\mathcal{P}$ of M is said to be a partition of M if each finite intersection of $\mathcal{P}$ is a PL-manifold, the geometric interiors of any two non-equal finite intersections are disjoint and for every non-empty finite intersection $P = P_0 \cap \dots \cap P_t$ of distinct elements $P_0, \dots, P_t \in \mathcal{P}$, $\dim P = \dim M - t$.

A decomposition $\mathcal{P}$ of M is said to be a partition on a PL-submanifold N of M if $\mathcal{P}|N$ is a partition of N (in this case we also say that $\mathcal{P}$ forms a partition on N or $\mathcal{P}$ restricted to N is a partition).

A subset P of M is said to be l -co-connected, $l \geq 0$, if P is $(\dim P - l)$ -connected (we assume that P is l -co-connected for every $l \geq \dim P + 1$). A partition $\mathcal{P}$ is said to be l -co-connected if every finite intersection of $\mathcal{P}$ is l -co-connected.

A map $f : X \rightarrow M$ is said to be in general position with a triangulation of M if for every simplex Δ of the triangulation $\dim f^{-1}(\Delta) \leq \dim X + \dim \Delta - \dim M$. Every map from X to M can be arbitrarily closely approximated by a general position map. Moreover,

let F be closed in X and a map $f : X \longrightarrow M$ restricted to F be in general position. Then f can be arbitrarily closely approximated by a general position map $g : X \longrightarrow M$ such that g coincides with f on F and $g(X \setminus F) \subset \text{Int}M$. A map $f : X \longrightarrow M$ is said to be in general position with a decomposition of M if it is in general position with a triangulation underlying this decomposition (a triangulation of M for which the elements of the decomposition are subcomplexes).

By an n -dimensional cube B^n we mean a set of the form $B^n = \{(x_1, \dots, x_n) : -r_i \leq x_i \leq r_i, r_i > 0, i = 1, \dots, n\}$ in the Euclidean space R^n . Thus we always assume that the origin O is at the center of B^n . Considering the product $B^n \times B^m$ of two cubes we identify B^n and B^m with $B^n \times O$ and $O \times B^m$ respectively.

Let $\mathcal{A}$ and $\mathcal{B}$ be collections of subsets of a set X and let $C \subset X$. We denote $\text{st}(C, \mathcal{A}) = \cup\{A : A \in \mathcal{A}, A \cap C \neq \emptyset\}$ and $\text{st}(\mathcal{A}, \mathcal{B}) = \{\text{st}(A, \mathcal{B}) : A \in \mathcal{A}\}$.

2.2 Elementary properties of partitions

If $\mathcal{P}$ is a partition of a triangulated manifold M then $\mathcal{P}|V$ is a partition of V for every open $V \subset M$. Let $\mathcal{P}$ be a decomposition of a triangulated manifold M and let $\mathcal{V}$ be an open cover M such that $\mathcal{P}|V$ is a partition of V for every $V \in \mathcal{V}$ then $\mathcal{P}$ is a partition of M .

Assume that M is a triangulated space. One can show that $M \times (0, 1)$ or $M \times [0, 1]$ is a PL-manifold if and only if M is a PL-manifold. This implies that for a triangulated manifold N , the product $M \times N$ is a PL-manifold if and only if M is a PL-manifold. Thus if M and N are triangulated manifolds and $\mathcal{P}$ is a decomposition of M then the decomposition $\mathcal{P} \times N = \{P \times N : P \in \mathcal{P}\}$ is a partition of $M \times N$ if and only if $\mathcal{P}$ is a partition of M .

The following properties are proved in [2] for compact manifolds however their proof also applies for the non-compact case.

Let $\mathcal{P}$ be a partition of a triangulated manifold M , let $\mathcal{P} = \cup\{\mathcal{P}_i : i = 1, 2, \dots\}$ be a splitting of $\mathcal{P}$ into disjoint subfamilies $\mathcal{P}_i$ and let $Q_i = \cup\{P : P \in \mathcal{P}_i\}$. Then $\mathcal{Q} = \{Q_1, Q_2, \dots\}$ is a partition of M (see 1.1.5 of [2]). In particular, any union of elements of a partition is a PL-manifold.

Let $\mathcal{P}$ be a partition of a triangulated manifold M . Then for every finite intersection P of $\mathcal{P}$, $P \cap \partial M \subset \partial P$ (see 1.1.9 of [2]).

Let M be an m -dimensional triangulated manifold, $\mathcal{P}$ a partition of M and $\mathcal{Q}$ a decomposition of M such that for every finite intersection P of $\mathcal{P}$, $\mathcal{Q}|P$ is a partition of P . Then $\mathcal{Q}$ is a partition of M (see 1.1.11 of [2]). In particular, if $M = M_1 \cup M_2$ is a decomposition of M into two m -dimensional PL-submanifolds M_1 and M_2 such that $N = M_1 \cap M_2$ is an $(m - 1)$ -dimensional PL-submanifold of both ∂M_1 and ∂M_2 and, $\mathcal{Q}$ is a decomposition of M such that $\mathcal{Q}|M_1$, $\mathcal{Q}|M_2$ and $\mathcal{Q}|N$ are partitions then $\mathcal{Q}$ is a partition of M .

2.3 A Matching of partitions

Let $\mathcal{P}_1$ and $\mathcal{P}_2$ be partitions of triangulated manifolds M_1 and M_2 respectively. A one-to-one correspondence $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ is said to be a matching of $\mathcal{P}_1$ and $\mathcal{P}_2$ if it induces a one-to-one correspondence between non-empty intersections. This means that for every finite intersection $P = P_0 \cap \dots \cap P_t$ of distinct elements $P_0, \dots, P_t \in \mathcal{P}_1$, $\mu(P) = \mu(P_0) \cap \dots \cap \mu(P_t) = \emptyset$ if and only if $P = \emptyset$.

Assume that $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ is a matching of partitions $\mathcal{P}_1$ and $\mathcal{P}_2$ such that $\mathcal{P}_2$ is l -co-connected. Let F be a closed subset of X such that $\dim X \leq m_i - l + 1$, $m_i = \dim M_i$ $i = 1, 2$, and let $f_i : F \longrightarrow M_i$ be maps such that $f_1^{-1}(P) = f_2^{-1}(\mu(P))$ for every finite intersection P of $\mathcal{P}_1$. Assume that a map $f'_1 : X \longrightarrow M_1$ extends f_1 such that f'_1 is in general position with $\mathcal{P}_1$. Then there is a map $f'_2 : X \longrightarrow M_2$ such that f'_2 extends f_2 and $(f'_1)^{-1}(P) = (f'_2)^{-1}(\mu(P))$ for every finite intersection P of $\mathcal{P}_1$.

The required extension of f_2 can be constructed by induction on the co-dimension $t = m_2, m_2 - 1, m_2 - 2, \dots, 0$ of the intersections of $\mathcal{P}_2$. Assume that we already extended f_2 to f'_2 on X_t = the union of the preimages of all the intersections of $\mathcal{P}_2$ of $\dim \leq m_2 - t$ such that f'_2 on X_t satisfies the required properties and let an intersection P of $\mathcal{P}_2$ be of $\dim = m_2 - t + 1$. Then for $X_P = (f'_1)^{-1}(\mu^{-1}(P))$, $\dim X_P \leq \dim X + \dim \mu^{-1}(P) - m_1 = \dim X + \dim P - m_2$. Since P is $(\dim P - l)$ -connected and $\dim P - l \geq \dim P - m_2 + \dim X - 1 \geq \dim X_P - 1$, f'_2 restricted to $X_P \cap X_t$ can be extended over X_P such that $f'_2(X_P \setminus X_t) \subset \text{Int}P$.

Thus we can extend f_2 to a map f'_2 with the required properties. This extension will be called a transfer of the extension f'_1 via the matching μ .

Assume that $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ is a matching of partitions on manifolds M_1 and M_2 respectively such that $\mathcal{P}_i$ is l_i -co-connected and $m_1 - l_1 = m_2 - l_2$ where $m_i = \dim M_i$. Let $n \leq m_1 - l_1$ and let $P_1 \subset P'_1$ be subsets of M_1 such that P_1, P'_1 are unions of elements of $\mathcal{P}_1$ and the inclusion $P_1 \subset P'_1$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n$. Then the inclusion $P_2 = \mu(P_1) \subset P'_2 = \mu(P'_1)$ also induces the zero-homomorphism of the homotopy groups in $\dim \leq n$. Indeed, take a map $f_2 : S^p \longrightarrow P_2$, $p \leq n$ from a p -dimensional sphere S^p into P_2 . Since P_2 is a manifold we can homotope f_2 inside P_2 to a general position map. Thus we assume that f_2 is a general position map and transfer this map via the matching μ to $f_1 : S^p \longrightarrow P_1$. Extend f_1 to a general position map $f'_1 : B^{p+1} \longrightarrow P'_1$ of an $(p+1)$ -dimensional ball B^{p+1} such that $S^p = \partial B^{p+1}$ and once again transfer this extension to an extension $f'_2 : B^{p+1} \longrightarrow P'_2$ of f_2 . Thus f_2 is null-homotopic in P'_2 .

2.4 A black hole modification

Let B^q, B^{m-q}, B_*^{m-q} and $B^m = B^q \times B^{m-q}$ be cubes with the dimensions indicated by the superscripts such that B_*^{m-q} is contained in $\text{Int}B^{m-q}$ and the centers of B_*^{m-q} and B^{m-q} coincide (recall that we assume that the centers of the cubes are located at O and we identify B^q and B^{m-q} with the subsets $B^q \times O$ and $O \times B^{m-q}$ of B^m respectively). Denote $T = B^q \times (B^{m-q} \setminus \text{Int}B_*^{m-q})$ and $S = \partial B^{m-q}$. Note that S is an $(m - q - 1)$ -dimensional

sphere in ∂T and T is a subset of B^m which is PL-homeomorphic to the product $B^{q+1} \times S$ where S is identified with $O \times S$.

Assume that B^m is PL-embedded in $\text{Int}M$ of a triangulated m -dimensional manifold M . Denote $L = M \setminus \text{Int}T$ and assume that F_L is a PL-subcomplex of L and $\mathcal{P}_L$ is a decomposition of L such that $S \subset F_L$ and $\mathcal{P}_L$ restricted to $L \setminus F_L$ and to $\partial T \setminus F_L$ are partitions.

Fix a triangulation $\mathcal{T}$ of M that underlies S , F_L and the elements of $\mathcal{P}_L$ and denote by $\mathcal{A}$ the collection of subcomplexes of L with respect to this triangulation. We are going to extend each A in $\mathcal{A}$ to a PL-subcomplex $A \subset A^\pi$ of M such that the following conditions are satisfied:

- $\mathcal{P}_L^\pi = \{P_L^\pi : P_L \in \mathcal{P}_L\}$ is a decomposition of M which forms a partition on $M \setminus F_L^\pi$;
- $F_L^\pi \cap L = F_L$ and $\mathcal{P}_L^\pi$ restricted to $L \setminus F_L^\pi$ coincides with $\mathcal{P}_L$ restricted to $L \setminus F_L$;
- the correspondence between $\mathcal{P}_L$ and $\mathcal{P}_L^\pi$ defined by $P_L \longrightarrow P_L^\pi$ induces a matching of partitions when $\mathcal{P}_L$ and $\mathcal{P}_L^\pi$ are restricted to $L \setminus F_L$ and $M \setminus F_L^\pi$ respectively;
- for every intersection $P_L = P_L^0 \cap \dots \cap P_L^j$ of elements $P_L^0, \dots, P_L^j$ of $\mathcal{P}_L$, $P_L^\pi \setminus F_L^\pi = ((P_L^0)^\pi \cap \dots \cap (P_L^j)^\pi) \setminus F_L^\pi$ and $P_L \setminus F_L$ is a (strong) deformation retract of $P_L^\pi \setminus F_L^\pi$;
- $\dim(F_L^\pi \cap T) \setminus S \leq \dim(F_L \cap \partial T) \setminus S + 1$;
- and finally $F_L^\pi = S$ if $F_L = S$.

Note that $T \setminus S$ can be represented as the product $(\partial T \setminus S) \times [0, 1)$ where $\partial T \setminus S$ corresponds to $(\partial T \setminus S) \times 0$. One of the ways to construct A^π is to find a retraction $\pi : T \setminus S \longrightarrow \partial T \setminus S$ which is topologically equivalent to the projection of $(\partial T \setminus S) \times [0, 1)$ to $(\partial T \setminus S) \times 0$ such that for A^π defined by $A^\pi = A$ if $A \cap \partial T = \emptyset$ and $A^\pi = A \cup S \cup \pi^{-1}(A \cap \partial T)$ otherwise we have that A^π is a PL-subcomplex of M for every $A \in \mathcal{A}$. Such a retraction π can be constructed however the author could not find an elementary and direct argument for this. Therefore for the sake of elementariness (and at expense of a geometric transparency) we adopt a slightly different approach for constructing A^π which is presented in 4.1. Note that all the properties of $\mathcal{P}_L^\pi$ and F_L^π that will be used in the sequel are described above. Thus for understanding the rest of the paper the reader may assume that the construction is carried out on base of an appropriate retraction π .

We will call the decomposition $\mathcal{P}_L^\pi$ a black hole modification of $\mathcal{P}_L$ with the sphere S as the black hole of the modification.

2.5 Improving connectivity of an intersection

Assume that $m \geq 2q + 1$ and $l = m - q + 2$, M is a triangulated m -dimensional $(q - 1)$ -connected ($= (l - 1)$ -co-connected) manifold and F is a PL-subcomplex of M lying in $\text{Int}M$. Let $\mathcal{P}$ be a decomposition of M which forms an l -co-connected partition on $U = M \setminus F$ and let $P = P_0 \cap P_1 \cap \dots \cap P_t$, $0 \leq t \leq m - l + 1$ be an intersection of distinct elements $P_0, \dots, P_t$ of $\mathcal{P}$ such that $P \cap U \neq \emptyset$. Then $P \cap U$ is an $(m - t)$ -dimensional and $(q - t - 2)$ -connected ($= l$ -co-connected) manifold.

Assume that the intersections of $\mathcal{P}|U$ of dimension $> m - t$ are $(l - 1)$ -co-connected. Let us show how to improve the connectivity of $P \cap U$ preserving the level of connectivity

of the rest of the intersections.

Let $f : S_P \longrightarrow \text{Int}(P \cap U)$ be a PL-embedding of a triangulated $(q-t-1)$ -dimensional sphere S_P such that f is not null-homotopic in $P \cap U$.

Consider a t -dimensional simplex Δ with vertices $V(\Delta) = \{v_0, \dots, v_t\}$. Let Δ' be a (sub)simplex of Δ lying in $\partial\Delta$ and let $V(\Delta') \subset V(\Delta)$ be the set of vertices of Δ' . Denote $P(\Delta') = \cap\{P_i : v_i \in V(\Delta) \setminus V(\Delta')\}$. Note that $P(\emptyset) = P$ and $P(\Delta') \subset P(\Delta'')$ if $\Delta' \subset \Delta''$.

For every simplex Δ' of $\partial\Delta$ we will define by induction on $\dim \Delta'$ a PL-embedding $f_{\Delta'} : S_P * \Delta' \longrightarrow P(\Delta') \cap U$ such that $f_{\Delta''}$ extends $f_{\Delta'}$ if $\Delta' \subset \Delta''$. Set $f_\emptyset = f$ and we are done for $t = 0$. Let $t > 0$. Take a simplex Δ' of $\dim = i + 1 \leq t - 1$ and assume that $f_{\Delta'}$ is defined for all Δ' of $\dim \leq i$. Then the maps $f_{\Delta''}$ for $\Delta'' \subset \partial\Delta'$ define the corresponding PL-embedding $f_{\partial\Delta'} : S_P * \partial\Delta' \longrightarrow \partial(P(\Delta') \cap U)$. Note that $S_P * \partial\Delta'$ is a $(q-t+i)$ -dimensional sphere and $P(\Delta') \cap U$ is $(m-t+i+2)$ -dimensional and $(q-t+i+1)$ -connected ($= (l-1)$ -co-connected) and $\dim(P(\Delta') \cap U) = m-t+i+2 \geq 2q+1-t+i+2 \geq 2(q-t+i+1)+1+(t-i-2) \geq 2(q-t+i+1)+1$ since $m \geq 2q+1$ and $t-i-2 \geq 0$. Then $f_{\partial\Delta'}$ can be extended to a PL-embedding $f_{\Delta'} : S_P * \Delta' \longrightarrow P(\Delta') \cap U$ such that $f_{\Delta'}$ sends $\text{Int}(S_P * \Delta')$ to $\text{Int}(P(\Delta') \cap U)$.

The maps $f_{\Delta'}$ define the PL-embedding $f_{\partial\Delta}$ of $S_P * \partial\Delta$ to $\text{Int}((P_0 \cup \dots \cup P_t) \cap U)$. Identify the $(q-1)$ -dimensional sphere $S^{q-1} = S_P * \partial\Delta$ with its image under $f_{\partial\Delta}$ and, by 4.2, extend the embedding of S^{q-1} to a PL-embedding of a q -dimensional cube B^q such that $S^{q-1} = \partial B^q$ and the embedding of B^q extends to a PL-embedding of an m -dimensional cube $B^m = B^q \times B^{m-q} \subset \text{Int}M$ having the following properties: $P' \cap B^m = (B^q \cap P') \times B^{m-q}$ for every $P' \in \mathcal{P}$; $F \cap B^m = (B^q \cap F) \times B^{m-q}$; $\mathcal{P}$ restricted to $N \cap U$ and $\partial B^m \cap U$ are partitions where $N = M \setminus \text{Int}B^m$ and for every finite intersection P' of $\mathcal{P}$ such that $P' \cap U \neq \emptyset$ we have that $P' \cap N \cap U \neq \emptyset$ and $P' \cap N \cap U$ is a deformation retract of $(P' \cap U) \setminus \text{Int}B^q$. Note that for every finite intersection P' of $\mathcal{P}$ that intersects U , $(\dim(P' \cap U) - (l-1) + 1) + q + 1 = \dim(P' \cap U) - m + 2q + 1 \leq \dim(P' \cap U)$ and therefore the inclusion $(P' \cap U) \setminus \text{Int}B^q \subset P' \cap U$ induces an isomorphism of the homotopy groups in $\dim \leq \dim(P' \cap U) - (l-1)$ ($=$ in co-dimensions $\geq l-1$). Also note that since $F \cap \partial B^q = \emptyset$ we have $F \cap \partial B^m = (F \cap \text{Int}B^q) \times \partial B^{m-q}$ and therefore $\dim F \cap \partial B^m \leq \dim F - 1$.

Let B_*^{m-q} be a cube lying in the geometric interior of B^{m-q} such that the centers of B_*^{m-q} and B^{m-q} coincide and $\partial B^q \times B_*^{m-q} \subset \text{Int}((P_0 \cup \dots \cup P_t) \cap U)$.

Use the notations of 2.4. Define $F_L = (N \cap F) \cup S$ where $S = \partial B^{m-q}$. Consider B^q as the join $O * \partial B^q$ where O is the center of B^q and define a decomposition $\mathcal{P}_L$ of L as the collection of the sets $\mathcal{P}_L = \{P'_L : P' \in \mathcal{P}\}$ where $P'_L = P' \cap N$ if $P' \in \mathcal{P}$ and $P' \neq P_0, \dots, P_t$ and $P'_L = (P' \cap N) \cup ((O * (P' \cap \partial B^q)) \times B_*^{m-q})$ for $P' = P_i$, $0 \leq i \leq t$. Note that $\mathcal{P}_L$ restricted to N coincides with $\mathcal{P}$ restricted to N . It is easy to see that $\mathcal{P}_L$ restricted to $B^q \times B_*^{m-q}$, $\partial B^q \times (B^{m-q} \setminus \text{Int}B_*^{m-q})$, $\partial B^q \times B_*^{m-q}$, $B^q \times \partial B_*^{m-q}$ and $\partial B^q \times \partial B_*^{m-q}$ are partitions. Then, by 2.2, $\mathcal{P}_L$ is a partition on $L \setminus (N \cap F)$ and $\partial T \setminus (N \cap F)$ and hence $\mathcal{P}_L$ is a partition on $L \setminus F_L$ and $\partial T \setminus F_L$. For a finite intersection $P' = P'_0 \cap \dots \cap P'_j$ of $P'_0, \dots, P'_j \in \mathcal{P}$ such that $P' \cap U \neq \emptyset$ denote $P'_L = (P'_0)_L \cap \dots \cap (P'_j)_L$.

Now apply the black hole modification to $\mathcal{P}_L$ and F_L with S as the black hole of the modification. Note that since S lies in ∂N then, by 2.2, for every finite intersection P' of $\mathcal{P}$, $P' \cap U \cap N \cap S \subset \partial(P' \cap U \cap N)$ and hence the inclusion $(P' \cap U \cap N) \setminus S \subset P' \cap U \cap N$ induces

an isomorphism of the homotopy groups in all dimensions. Recall that $f(S_P) \subset P \cap U \cap N$. By 2.4 and the construction of $\mathcal{P}_L$ one can easily verify that for every finite intersection P' of $\mathcal{P}$ such that $P' \cap U \neq \emptyset$ and $P \neq P'$, $(P' \cap U \cap N) \setminus S$ is a deformation retract of $(P'_L)^\pi \setminus F_L^\pi$ and for the intersection P the inclusion of $(P \cap U \cap N) \setminus S$ into $P_L^\pi \setminus F_L^\pi$ induces an isomorphism of the homotopy groups in dimensions $< q - t - 1$ and an epimorphism in $\dim = q - t - 1$ such that the element of the homotopy group represented by the map f is in the kernel of this epimorphism.

Thus for every intersection P' of $\mathcal{P}$ such that $P' \cap U \neq \emptyset$ the connectivity of $(P'_L)^\pi \setminus F_L^\pi$ fits the connectivity of $P' \setminus F$ for $P' \neq P$ and we contributed toward improving the connectivity of $P \setminus F$ in its modification $P_L^\pi \setminus F_L^\pi$.

Denote $F^\pi = F_L^\pi$, $(P')^\pi = (P'_L)^\pi$ for a finite intersection P' of $\mathcal{P}$ and let $\mathcal{P}^\pi = \{(P')^\pi : P' \in \mathcal{P}\} = \mathcal{P}_L^\pi$. It is easy to see that the one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}_L$ defined by $P' \longrightarrow (P')_L, P' \in \mathcal{P}$ becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}_L$ are restricted to $M \setminus F$ and $L \setminus F_L$ respectively. Then by 2.4 the one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}^\pi$ defined by $P' \longrightarrow (P')^\pi, P' \in \mathcal{P}$ becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}^\pi$ are restricted to $M \setminus F$ and $M \setminus F^\pi$ respectively. Note that $F^\pi \subset \text{Int}M$ and because $\dim(F \cap \partial B^m) \leq \dim F - 1$ we have $\dim F^\pi \leq \max\{\dim S, \dim F\} = \max\{m - q - 1, \dim F\}$.

Finally note that even if $F = \emptyset$ we have $F^\pi = S$ and we improve the connectivity of P at expense of an "irregular" behavior of $\mathcal{P}^\pi$ on the black hole S , in particular, we create additional intersections of $\mathcal{P}^\pi$ on S and the elements of $\mathcal{P}^\pi$ are no longer manifolds in M .

2.6 A discretization of maps' images

Assume that M be a triangulated manifold such that $\dim M \geq 2r + 1$ and $\mathcal{U}$ is an open cover of M .

Let $\mathcal{F}$ be countable collection of maps $f : S^r \longrightarrow M$ from a triangulated r -dimensional sphere S^r . Then there is a PL-subcomplex R of M of $\dim \leq r$ lying in $\text{Int}M$ such that every map f in $\mathcal{F}$ admits a $\mathcal{U}$ -close approximation $f' : S^r \longrightarrow \text{Int}M \setminus R$ such that f' is a PL-embedding and the images of the maps in $\mathcal{F}' = \{f' : f \in \mathcal{F}\}$ form a discrete family in $M \setminus R$. Moreover, given a decomposition $\mathcal{P}$ of M , the set R can be chosen so that R is nowhere dense on the finite intersections of $\mathcal{P}$, that is the intersection of R with every finite intersection of $\mathcal{P}$ is nowhere dense in that intersection.

Indeed, arrange $\mathcal{F}$ into a sequence $\mathcal{F} = \{f_1, f_2, \dots\}$ and take a sufficiently small triangulation $\mathcal{T}$ of M such that $\mathcal{T}$ underlies $\mathcal{P}$. Approximate each $f_i : S^{r-1} \longrightarrow M$ by a map $f_i'' : S^r \longrightarrow M$ such that $f_i''(S^r)$ is contained in R' = the r -skeleton of $\mathcal{T}$. Fix a metric d on M and approximate each f_i'' by an PL-embedding $f_i''' : S^r \longrightarrow M$ such that $f_i'''(S^r) \cap R = \emptyset$, $d(f_i'''(S^r), R') \leq 1/i$ and $f_i'''(S^r) \cap f_j'''(S^r) = \emptyset$ for $i \neq j$. Then the sets $f_i'''(S^r), i = 1, 2, \dots$ form a discrete family in $M \setminus R'$. Now take a PL-embedding $g : M \longrightarrow M$ such that $g(M)$ is a PL-subcomplex of M , $g(M) \subset \text{Int}M$, the restriction of g to R' is in general position with $\mathcal{T}$ and g is close to the identity map of M . It is clear that the construction can be carried out such that $f_i' = g \circ f_i'''$ is $\mathcal{U}$ -close to f_i . Then the maps f_i' and the set $R = g(R')$ will have the required properties.

In a similar way one can show that if $\mathcal{F}$ is a countable collection of maps $f : B^r \rightarrow M$ from a triangulated r -dimensional ball B^r such that for every f in $\mathcal{F}$, f restricted to ∂B^r is a PL-embedding and the images of the maps in $\{f|_{\partial B^r} : f \in \mathcal{F}\}$ form a discrete family in M then there is a PL-subcomplex R of M of $\dim \leq r$ lying in $\text{Int} M$ such that every map f in $\mathcal{F}$ admits a $\mathcal{U}$ -close approximation $f' : B^r \rightarrow M \setminus R$ such that f' is a PL-embedding, $f'|_{\partial B^r} = f|_{\partial B^r}$, $f'(\text{Int} B^r) \subset \text{Int} M$ and the images of the maps in $\mathcal{F}' = \{f' : f \in \mathcal{F}\}$ form a discrete family in $M \setminus R$. Moreover, given a decomposition $\mathcal{P}$ of M , the set R can be chosen so that R is nowhere dense on the finite intersections of $\mathcal{P}$.

Note that if M is $(r-1)$ -connected and R is a closed PL-presented subset of M of $\dim \leq r$ then $M' = M \setminus R$ is $(r-1)$ -connected. In addition, if $\mathcal{P}$ is a decomposition of M which is an l -co-connected partition on an open subset U of M with $l \geq r+2$ and R is nowhere dense on the finite intersections of $\mathcal{P}$ then $\mathcal{P}' = \{P \setminus R : P \in \mathcal{P}\}$ is a decomposition of M' which is an l -co-connected partition on $U' = U \setminus R$ and the one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}'$ defined by $P \rightarrow P \setminus R, P \in \mathcal{P}$ induces a matching of partitions when $\mathcal{P}$ and $\mathcal{P}'$ are restricted to U and U' respectively (the assumption that R is nowhere dense on the finite intersections of $\mathcal{P}$ is needed to guarantee that for every finite intersection P of $\mathcal{P}$, $(P \cap U) \setminus R \neq \emptyset$ provided $P \cap U \neq \emptyset$).

2.7 Improving connectivity of intersections simultaneously

Adopt the notations and the assumptions of 2.5. We are going to show how to improve simultaneously the connectivity of all the intersections of $\mathcal{P}|_U$ of $\dim = m-t$ to $(l-1)$ -co-connectivity.

For an intersection P of $\mathcal{P}$ with $\dim P \cap U = m-t$ choose a countable collection $\mathcal{F}_P$ of PL-embeddings $f : S_P \rightarrow \text{Int}(P \cap U)$ from a $(q-t-1)$ -dimensional sphere S_P that generate the $(q-t-1)$ -dimensional homotopy group of $P \cap U$. Since F is a PL-subcomplex of M we may assume that the images of the maps in $\mathcal{F}_P$ lie outside a neighborhood of F . Then by 2.6 one can find a PL-subcomplex R_P of M such that $R_P \subset \text{Int}(P \cap U)$ and $\dim R_P \leq q-t-1$ and assume that the images of the embeddings in $\mathcal{F}_P$ are contained in $\text{Int}((P \cap U) \setminus R_P)$ and form a discrete family in $M \setminus R_P$.

Recall that in 2.5 the elements of $\mathcal{P}$ that form the intersection P are enumerated according to the vertices of a sample simplex $\Delta = \{v_0, \dots, v_t\}$ (we fix such enumeration arbitrarily and independently for every intersection P of $\mathcal{P}$ with $\dim P \cap U = m-t$). Following 2.5 and using 2.6 we can define for every $\Delta' \subset \Delta$ a set $R_P^{\Delta'}$ and collections of maps $\mathcal{F}_P^{\Delta'} = \{f_{\Delta'} : f \in \mathcal{F}_P\}$ and $\mathcal{F}_P^{\partial \Delta'} = \{f_{\partial \Delta'} : f \in \mathcal{F}_P\}$ such that $R_P^{\Delta''} \subset R_P^{\Delta'}$ if $\Delta'' \subset \Delta'$, $R_P^{\Delta'}$ is a closed subset of M lying in $P(\Delta') \cap U$, the images of the maps of $\mathcal{F}_P^{\Delta'}$ are contained in $G_P^{\Delta'} = (P(\Delta') \cap U) \setminus R_P^{\Delta'}$ and form a discrete family in $M \setminus R_P^{\Delta'}$, and $R_P^{\Delta'} \setminus R_P^{\partial \Delta'}$ is a PL-subcomplex of $M \setminus R_P^{\partial \Delta'}$ lying in $(P(\Delta') \cap U) \setminus R_P^{\partial \Delta'}$ such that $\dim R_P^{\Delta'} \setminus R_P^{\partial \Delta'} \leq q-t+\dim \Delta'$ where $R_P^{\partial \Delta'} = \cup \{R_P^{\Delta''} : \Delta'' \subset \partial \Delta'\}$. Here we assume that $\Delta' = \emptyset$ is a simplex of Δ and $\mathcal{F}_P^\emptyset = \mathcal{F}_P$ and $R_P^\emptyset = R_P$ and we also assume that $P(\Delta) = M$.

Denote by $\mathcal{F}^{\partial \Delta}$ and $R^{\partial \Delta}$ the union of $\mathcal{F}_P^{\partial \Delta}$ and $R_P^{\partial \Delta}$ respectively over all the intersections P of $\mathcal{P}$ with $\dim P \cap U = m-t$. In order to carry out the construction described

above such that $R^{\partial\Delta}$ will be a closed PL-presented subset of M and the images of the maps in $\mathcal{F}^{\partial\Delta}$ will form a discrete family in $M \setminus R^{\partial\Delta}$, we need to take into account that the same intersection P' of $\mathcal{P}$ with $\dim P' \cap U > m - t$ can be involved in many (even countably many) intersections of $\dim = m - t$. It can be done as follows. For a finite intersection P' of $\mathcal{P}$ with $\dim P' \cap U > m - t$ denote by $R_{\partial P'}$, $R_{P'}$, $\mathcal{F}_{\partial P'}$ and $\mathcal{F}_{P'}$ the union of $R_P^{\partial\Delta'}$, $R_P^{\Delta'}$, $\mathcal{F}_P^{\partial\Delta'}$ and $\mathcal{F}_P^{\Delta'}$ respectively over all finite intersections P of $\mathcal{P}$ such that $\dim P \cap U = m - t$ and $\Delta' \subset \partial\Delta$ such that $P(\Delta') = P'$ and, let $G_{P'} = (P' \cap U) \setminus R_{P'}$. It is obvious that if $\dim P' \cap U = m - t + 1$ then $R_{\partial P'}$ is closed in M , the images of the maps in $\mathcal{F}_{\partial P'}$ are contained in $\partial(P' \cap U) \setminus R_{\partial P'}$ and form a discrete family in $M \setminus R_{\partial P'}$. Now assume that the last properties hold for a finite intersection P' of $\mathcal{P}$ with $\dim P' \cap U > m - t$. Then applying 2.6 to all the maps of $\mathcal{F}_{\partial P'}$ we can enlarge $R_{\partial P'}$ to a closed subset $R_{P'}$ of M contained in $P' \cap U$ and find extensions $\mathcal{F}_{P'}$ of the maps in $\mathcal{F}_{\partial P'}$ such that $\dim R_{P'} \leq \dim(P' \cap U) - l + 1 = \dim(P' \cap U) + q - m - 1$, $R_{P'} \setminus R_{\partial P'}$ is a PL-subcomplex of $M \setminus R_{\partial P'}$, the images of the extensions $\mathcal{F}_{P'}$ of the maps in $\mathcal{F}_{\partial P'}$ are contained in $G_{P'}$ and form a discrete family in $M \setminus R_{P'}$. Thus we can define $R_P^{\Delta'} = R_{P'}$ for every finite intersection P of $\mathcal{P}$ with $\dim P \cap U = m - t$ and $\Delta' \subset \partial\Delta$ such that $P(\Delta') = P'$. Then proceeding by induction on $\dim P' \cap U$ we construct for every intersection P of $\mathcal{P}$ with $\dim P \cap U = m - t$ and $\Delta' \subset \partial\Delta$ the collection of maps $\mathcal{F}_P^{\Delta'}$ and the set $R_P^{\Delta'}$ such that $R^{\partial\Delta}$ is a closed PL-presented subset of M contained in U such that $\dim R^{\partial\Delta} \leq q - 1$ and the images of the maps in $\mathcal{F}^{\partial\Delta}$ are contained in $(M \cap U) \setminus R^{\partial\Delta}$ and form a discrete family in $M \setminus R^{\partial\Delta}$. From the construction it is clear that $R^{\partial\Delta}$ is nowhere dense on the finite intersections of $\mathcal{P}$.

By 2.6 the set $R^{\partial\Delta}$ can be enlarged to a closed PL-presented subset R of M and each map $f_{\partial\Delta} : S^{q-1} \longrightarrow M \setminus R^{\partial\Delta}$ in $\mathcal{F}^{\partial\Delta}$ can be extended to a map $f'_\Delta : B^q \longrightarrow M \setminus R$ such that $\dim R \leq q$, R is nowhere dense on the finite intersections of $\mathcal{P}$ and the images of the maps f'_Δ form a discrete family in $M \setminus R$. Hence for every $f_{\partial\Delta} \in \mathcal{F}^{\partial\Delta}$ there is an open subset $Q(f_{\partial\Delta})$ of $M \setminus R$ containing the image of $f_{\partial\Delta}$ such that $\mathcal{Q} = \{Q(f_{\partial\Delta}) : f_{\partial\Delta} \in \mathcal{F}^{\partial\Delta}\}$ is a discrete family in $M \setminus R$ and $f_{\partial\Delta}$ is null-homotopic in $Q(f_{\partial\Delta})$. Then by 4.2 the black hole modification used in 2.5 and involving $f_{\partial\Delta}$ can be carried out inside $Q(f_{\partial\Delta})$.

Note that since R is a PL-presented set of $\dim \leq q$ and $m \geq 2q + 1$ then for every finite intersection P of $\mathcal{P}$ that intersects U we have that the inclusion $(P \cap U) \setminus R \subset P \cap U$ induces an isomorphism of the homotopy groups in $\dim \leq \dim(P \cap U) - (l - 1)$ (=co-dimensions $\geq l - 1$).

Thus after removing R from M , F and the elements of $\mathcal{P}$ the black hole modification 2.4 used in 2.5 can be applied independently for every map in $\mathcal{F}^{\partial\Delta}$ in order to modify $\mathcal{P}$ to a decomposition $\mathcal{P}'$ of $M' = M \setminus R$ and F to a set F' such that $\mathcal{P}'$ = the result of all the modifications $\mathcal{P}^\pi$ and F' = the result of all the modifications F^π . Then it is easy to derive from 2.5 that F' is a PL-subcomplex of M' of $\dim \leq \max\{m - q - 1, \dim F\}$ lying in $\text{Int} M'$, $\mathcal{P}'$ is a partition on $M' \setminus F'$ and for every intersection P of $\mathcal{P}$ with $P \cap U \neq \emptyset$, P is modified to P' (= the result of all the modifications P^π) such that $P' \cap (M' \setminus F')$ is l -co-connected and $P' \cap (M' \setminus F')$ is $(l - 1)$ -co-connected if $\dim P \cap U \geq m - t$.

Note that the natural one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}'$ defined by sending $P \in \mathcal{P}$ to its modification $P' \in \mathcal{P}'$ turns into a matching of partitions when $\mathcal{P}$ and $\mathcal{P}'$ are

restricted to $M \setminus F$ and $M' \setminus F'$.

It is clear that the images of the maps in $\mathcal{F}^{\partial\Delta}$ are contained in the elements of $\text{st}(\mathcal{P}, \mathcal{P})$. Assume that $\text{st}(\mathcal{P}, \mathcal{P})$ refines an open cover $\mathcal{W}$ of M such that for every $A \in \text{st}(\mathcal{P}, \mathcal{P})$ there is $W \in \mathcal{W}$ such that $A \subset W$ and the inclusion of A into W induces the zero-homomorphism of the homotopy groups in $\dim \leq q - 1$. By 2.6 the collection $\mathcal{Q}$ can be chosen so that $\mathcal{Q}$ refines $\mathcal{W}$. Then it is easy to see that $\mathcal{P}'$ refines $\text{st}(\mathcal{W}, \mathcal{W})$.

2.8 Improving the total connectivity of a partition

Let M be a triangulated $(q - 1)$ -connected m -dimensional manifold with $m \geq 2q + 1$ and let $l = m - q + 2$. Assume that F is a PL-subcomplex of M lying in $\text{Int}M$ and $\mathcal{P}$ is a decomposition of M such that $\dim F \leq m - q$ and $\mathcal{P}$ is an l -co-connected partition on $U = M \setminus F$.

Apply 2.7 to improve the connectivity of the elements of $\mathcal{P}|U$ to the $(l - 1)$ -connectivity, again apply 2.7 to the modified decomposition to improve the connectivity of the intersections of $\dim = m - 1$ (of the modified decomposition restricted to the complement of the modified F) to the $(l - 1)$ -connectivity and thus proceed on the dimension of the intersections $\geq l - 1$ until we modify M to an open subset $M' \subset M$, $\mathcal{P}$ to a decomposition $\mathcal{P}'$ of M' and F to a PL-subcomplex F' of M' lying in $\text{Int}M'$ such that $M \setminus M'$ is a closed PL-presented subset of M , $\dim(M \setminus M') \leq q$, $\dim F' \leq \max\{m - q - 1, \dim F\} \leq m - q$, $\mathcal{P}'$ is an $(l - 1)$ -co-connected partition on $U' = M' \setminus F'$ and $\mathcal{P}$ admits a natural one-to-one correspondence to $\mathcal{P}'$ which sends each element of $\mathcal{P}$ to its modification in $\mathcal{P}'$ and which becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}'$ are restricted to U and U' respectively.

2.9 Absorbing simplexes

Let M be a triangulated $(q - 1)$ -connected m -dimensional manifold with $m \geq 2q + 1$ and let $l = m - q + 1$. Assume that F is a PL-subcomplex of M lying in $\text{Int}M$ such that $U = M \setminus F$ is l -co-connected and $\dim F \leq m - q$ and assume that $\mathcal{P}$ is a decomposition of M such that $\mathcal{P}$ is an l -co-connected partition on U . Fix a triangulation $\mathcal{T}$ of M for which F is a subcomplex and let F' be the $(m - q - 1)$ -skeleton of F . We will show how to modify M to an open subset $M' \subset M$ and $\mathcal{P}$ to a decomposition $\mathcal{P}'$ of M' such that $M \setminus M'$ is a PL-subcomplex of M of $\dim \leq q$, $\mathcal{P}'$ restricted to $U' = M' \setminus F'$ is an l -co-connected partition and $\mathcal{P}$ admits a natural one-to-one correspondence to $\mathcal{P}'$ which becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}'$ restricted to U and U' respectively.

Assume that the triangulation $\mathcal{T}$ that we fixed in M is the second barycentric subdivision of a triangulation for which F and the elements of $\mathcal{P}$ are subcomplexes. Let $\mathcal{T}_F$ be the collection of all $(m - q)$ -dimensional simplexes lying in F , S_Δ the link of $\Delta \in \mathcal{T}_F$ with respect to $\mathcal{T}$, $Q = \text{Int}(\Delta * S_\Delta)$ for all Δ in $\mathcal{T}_F$ and $N = M \setminus (Q \cup F)$. Then for every $\Delta \in \mathcal{T}_F$ the link S_Δ of Δ is an $(q - 1)$ -dimensional sphere lying in $\text{Int}M$, for different simplexes Δ_1 and Δ_2 in $\mathcal{T}_F$ the joins $\Delta_1 * S_{\Delta_1}$ and $\Delta_2 * S_{\Delta_2}$ do not intersect on U , for every finite intersection P of $\mathcal{P}$ that intersects U and for every $\Delta \in \mathcal{T}_F$, $(P \cap U) \setminus \text{Int}(\Delta * S_\Delta)$ is

a deformation retract of $P \cap U$. Note that N is a manifold which is a deformation retract of U and therefore N is l -co-connected.

Fix $\Delta \in \mathcal{T}_F$ and let an open subset V_Δ of N be such that $S^{q-1} = S_\Delta \subset V_\Delta$ and the inclusion of S^{q-1} into V_Δ is null-homotopic in V_Δ . Then by 4.3 there are an element $P_\Delta \in \mathcal{P}$ and a PL-embedding of a cube $B^m = B^q \times B^{m-q}$ into $\text{Int}M$ such that $B^m \subset (\Delta * S_\Delta) \cup V_\Delta$, $\Delta = 0 \times B^{m-q} = F \cap B^m$, $P \cap B^m = (P \cap B^q) \times B^{m-q}$ for every $P \in \mathcal{P}$, $\partial B^q \times B^{m-q} \subset \text{Int}P_\Delta$, $\mathcal{P}$ restricted to $(M \setminus \text{Int}B^m) \cap U$ and $\partial B^m \cap U$ are partitions and, $(P \setminus \text{Int}B^m) \cap U$ is a deformation retract of $(P \setminus \text{Int}B^q) \cap U$ for every finite intersection P of $\mathcal{P}$.

Let us briefly describe the general idea of the construction in 4.3. The inclusion of S^{q-1} can be extended to a PL-embedding of a q -dimensional cube $B_\#^q$ into V_Δ such that $S^{q-1} = \partial B_\#^q$ and the embedding of $B_\#^q$ extends to a PL-embedding of an m -dimensional cube $B_\#^m = B_\#^q \times B_\#^{m-q}$ into V_Δ such that $\partial B_\#^q \times B_\#^{m-q} \subset B_\#^m \cap \partial(\Delta * S_\Delta)$ and $P \cap B_\#^m = (P \cap B_\#^q) \times B_\#^{m-q}$ for every $P \in \mathcal{P}$. Let $P_\Delta \in \mathcal{P}$ be such that $\text{Int}(P_\Delta \cap B_\#^q) \neq \emptyset$ and let a q -dimensional cube $\sigma_\#$ be PL-embedded in $\text{Int}(P_\Delta \cap B_\#^q)$. Then the cube $B^m = (\Delta * S_\Delta) \cup ((B_\#^q \setminus \text{Int}\sigma_\#) \times B_\#^{m-q})$ can be represented as the product $B^m = B^q \times B^{m-q}$ with the required properties. See 4.3 for details.

Let B_*^{m-q} be a cube lying in the geometric interior of B^{m-q} . Use the notations of 2.4. Define $F_L = F \setminus \text{Int}B^m$, $P_L = P \setminus \text{Int}B^m$ if $P \neq P_\Delta$ and $P \in \mathcal{P}$, $(P_\Delta)_L = (P_\Delta \setminus \text{Int}B^m) \cup (B^q \times B_*^{m-q})$ and define $\mathcal{P}_L = \{P_L : P \in \mathcal{P}\}$. Then $S = \partial B^{m-q} = \partial \Delta \subset F_L$ and it is easy to see that $\mathcal{P}_L$ is an l -co-connected partition on $L \setminus F_L$ and $\mathcal{P}_L$ is a partition on $\partial T \setminus F_L$. Now apply the black hole modification with the black hole $S = \partial \Delta$ and denote $F^\pi = F_L^\pi$, $P^\pi = P_L^\pi$ for $P \in \mathcal{P}$ and $\mathcal{P}^\pi = \{P^\pi : P \in \mathcal{P}\} = \mathcal{P}_L^\pi$. Note that $F_L^\pi = F_L$ since $F \cap \partial B^m = \partial \Delta$. Then $\mathcal{P}^\pi$ is a decomposition of M which is an l -co-connected partition on the manifold $M \setminus F^\pi = (M \setminus F) \cup \text{Int}\Delta$ in which we have absorbed the geometric interior of Δ in the modification of P_Δ . Note that the one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}_L$ defined by $P \longrightarrow P_L$, $P \in \mathcal{P}$ becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}_L$ are restricted to $M \setminus F$ and $L \setminus F_L$ respectively. Then by 2.4 the one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}^\pi$ defined by $P \longrightarrow P^\pi$, $P \in \mathcal{P}$ becomes a matching of partitions when $\mathcal{P}$ and $\mathcal{P}^\pi$ are restricted to $M \setminus F$ and $M \setminus F^\pi$ respectively.

Note that each S_Δ can be contracted to a point in N outside some neighborhood of F in M . Then by 2.6 there is a PL-subcomplex R of M contained in $\text{Int}N$ and nowhere dense on the finite intersections of $\mathcal{P}$ with $\dim R \leq q$ such that the collection $\mathcal{V}$ of the sets $V_\Delta, \Delta \in \mathcal{T}_F$ can be chosen to be discrete in $N \setminus R$. Then replacing M by $M' = M \setminus R$ and removing R the elements of $\mathcal{P}$ we can perform all the black hole modifications independently for all $\Delta \in \mathcal{T}_F$ and get from all the modifications $\mathcal{P}^\pi$ the corresponding decomposition $\mathcal{P}'$ of M' with the required properties. It is clear that the natural one-to-one correspondence between $\mathcal{P}$ and $\mathcal{P}'$ defined by sending each element of $\mathcal{P}$ to its modification becomes a matching of partitions when the decompositions are restricted to $M \setminus F$ and $M' \setminus F'$.

Let $\mathcal{W}$ be an open cover of M such that $\text{st}(\mathcal{P}, \mathcal{P})$ refines $\mathcal{W}$ and assume that for every $S_\Delta, \Delta \in \mathcal{T}_F$ there is $W_\Delta \in \mathcal{W}$ such that $S_\Delta \subset W_\Delta$ and S_Δ can be contracted to a point in $W_\Delta \cap N$. Then the collection $\mathcal{V}$ can be chosen so that $\mathcal{V}$ refines $\mathcal{W}$ and this implies

that $\mathcal{P}'$ refines $\text{st}(\mathcal{W}, \mathcal{W})$.

2.10 Improving connectivity via a matching

Let M_i be an l_i -co-connected triangulated manifolds such that $m_i = \dim M_i \geq 2(m_i - l_i) + 3$, $i = 1, 2$ and $m_1 - l_1 = m_2 - l_2$. Suppose that $\mathcal{P}_1$ and $\mathcal{P}_2$ are partitions of M_1 and M_2 respectively such that there is a matching between $\mathcal{P}_1$ and $\mathcal{P}_2$, and $\mathcal{P}_1$ is l_1 -co-connected. We will show how to modify M_2 to M'_2 and $\mathcal{P}_2$ to $\mathcal{P}'_2$ such that M'_2 is an open l_2 -co-connected submanifold of M_2 , $M_2 \setminus M'_2$ is a PL-presented closed subset of M_2 , $\dim(M_2 \setminus M'_2) \leq l_2 - 2$ and $\mathcal{P}'_2$ is an l_2 -co-connected partition of M'_2 which admits a natural matching to $\mathcal{P}_2$ defined by sending each element of $\mathcal{P}_2$ to its modification in $\mathcal{P}'_2$.

By 2.8 modify M_2 to $M_2^0 \subset M_2$, $\mathcal{P}_2$ to $\mathcal{P}_2^0$ and construct a subset F^0 of M_2^0 such that $M_2 \setminus M_2^0$ is a PL-subcomplex of M_2 , $\dim(M_2 \setminus M_2^0) \leq 1$, F^0 is a PL-subcomplex of M_2^0 , $\dim F^0 \leq m_2 - 2$, $\mathcal{P}_2^0$ is a decomposition of M_2^0 such that $\mathcal{P}_2^0$ restricted to $M_2^0 \setminus F^0$ is an m_2 -co-connected partition that admits a matching to $\mathcal{P}_2$ and therefore to $\mathcal{P}_1$.

Now assume that for $0 \leq t < m_2 - l_2$ we have constructed M_2^{2t} , $\mathcal{P}_2^{2t}$ and F^{2t} such that that $M_2 \setminus M_2^{2t}$ is a PL-presented closed subset of M_2 , $\dim(M_2 \setminus M_2^{2t}) \leq t + 1$, F^{2t} is a PL-subcomplex of M_2^{2t} , $\dim F^{2t} \leq m_2 - t - 2$, $\mathcal{P}_2^{2t}$ is a decomposition of M_2^{2t} such that $\mathcal{P}_2^{2t}$ restricted to $M_2^{2t} \setminus F^{2t}$ is an $(m_2 - t)$ -co-connected partition that admits a matching to $\mathcal{P}_2$ and therefore to $\mathcal{P}_1$. Proceed to $t + 1$ as follows.

By 2.8 modify M_2^{2t} to $M_2^{2t+1} \subset M_2^{2t}$, $\mathcal{P}_2^{2t}$ to $\mathcal{P}_2^{2t+1}$, F^{2t} to F^{2t+1} such that $M_2^{2t} \setminus M_2^{2t+1}$ is a PL-subcomplex of M_2^{2t} , $\dim(M_2^{2t} \setminus M_2^{2t+1}) \leq t + 2$, F^{2t+1} is a PL-subcomplex of M_2^{2t+1} , $\dim F^{2t+1} \leq m_2 - t - 2$, $\mathcal{P}_2^{2t+1}$ is a decomposition of M_2^{2t+1} , $\mathcal{P}_2^{2t+1}$ restricted to $M_2^{2t+1} \setminus F^{2t+1}$ is an $(m_2 - t - 1)$ -co-connected partition that admits a matching to $\mathcal{P}_2^{2t}$ restricted to $M_2^{2t} \setminus F^{2t}$ and therefore to $\mathcal{P}_2$ and $\mathcal{P}_1$.

Then since $\mathcal{P}_1$ is l_1 -co-connected on the l_1 -co-connected manifold M_1 we conclude by 2.3 that $M_2^{2t+1} \setminus F^{2t+1}$ is $(m_2 - t - 1)$ -co-connected. Therefore, by 2.9, M_2^{2t+1} , F^{2t+1} and $\mathcal{P}_2^{2t+1}$ can be modified to M_2^{2t+2} , F^{2t+2} and $\mathcal{P}_2^{2t+2}$ respectively such that $M_2^{2t+2} \subset M_2^{2t+1}$, $M_2^{2t+1} \setminus M_2^{2t+2}$ is a PL-subcomplex of M_2^{2t+1} , $\dim(M_2^{2t+1} \setminus M_2^{2t+2}) \leq t + 2$, F^{2t+2} is a PL-subcomplex of M_2^{2t+2} , $\dim F^{2t+2} \leq m_2 - t - 3$, $\mathcal{P}_2^{2t+2}$ is a decomposition of M_2^{2t+2} , $\mathcal{P}_2^{2t+2}$ restricted to $M_2^{2t+2} \setminus F^{2t+2}$ is an $(m_2 - t - 1)$ -co-connected partition that admits a matching to $\mathcal{P}_2^{2t+1}$ restricted to $M_2^{2t+1} \setminus F^{2t+1}$ and therefore to $\mathcal{P}_2$ and $\mathcal{P}_1$.

Then for $t = m_2 - l_2$, $M'_2 = M_2^{2t} \setminus F^{2t}$ and $\mathcal{P}'_2 = \mathcal{P}_2^{2t}$ restricted to M'_2 will have the required properties (note that $m_2 - l_2 + 1 \leq l_2 - 2$ and therefore $\dim(M_2 \setminus M'_2) \leq l_2 - 2$).

2.11 Moving to a rational position

Let M be a triangulated manifold with the rational structure determined by a triangulation $\mathcal{T}$ (see Introduction). The PL-notions defined in 2.1 can be translated to the corresponding rational notions by referring to the rational structure instead of the PL-structure. For example, $P \subset M$ is a rational subspace if $P \subset M$ is rationally embedded in M , P is a rational subcomplex if there is a rational triangulation of M for which P

is a subcomplex, $\mathcal{P}$ is a rational decomposition of M if the elements of $\mathcal{P}$ are rational subcomplexes and so on.

Assume that $\mathcal{T}'$ is a (not necessarily rational) triangulation of M . We will show that the identity map of M can be arbitrarily closely approximated by a PL-homeomorphism $f : M \rightarrow M$ such that for every $\Delta' \in \mathcal{T}'$, $f(\Delta')$ is a rational subcomplex of M .

Embed M into a Hilbert space by a map which is linear on every simplex of $\mathcal{T}$ and refer to this Hilbert space when properties of linearity are used. Let $\mathcal{T}''$ be a triangulation of M such that $\mathcal{T}''$ is a subdivision of both $\mathcal{T}$ and $\mathcal{T}'$, and the simplexes of $\mathcal{T}''$ are linear. Every vertex v of $\mathcal{T}''$ approximate by a rational point p_v (= a point with rational barycentric coordinates with respect to $\mathcal{T}$) such that for every simplex Δ of $\mathcal{T}$, $p_v \in \Delta$ if and only if $v \in \Delta$. The approximation of the vertices of $\mathcal{T}''$ can be extended to the PL-map $f : M \rightarrow M$ sending each vertex v of $\mathcal{T}''$ to p_v such that f is linear on each simplex of $\mathcal{T}''$. If p_v is sufficiently close to v for every vertex $v \in \mathcal{T}''$ then the map f is a PL-homeomorphism that can be chosen to be arbitrarily close to the identity map. Clearly f sends every simplex of $\mathcal{T}''$ to a rational simplex and therefore f sends every simplex of $\mathcal{T}'$ to a rational subcomplex of M .

Let M be an open subset of a space Y and d a metric on Y . We can assume that the homeomorphism $f : M \rightarrow M$ constructed above satisfies $d(y, f(y)) \leq d(y, Y \setminus M)$ for every $y \in M$. Then f extends to the homeomorphism $g : Y \rightarrow Y$ such that $g(y) = y$ if $y \in Y \setminus M$ and $g(y) = f(y)$ if $y \in M$. Now assume that R is a PL-presented subset of M and let $R_1 \subset R_2 \subset \dots \subset R_n$ be closed subsets of M such that $R = R_n$, R_1 is a PL-subcomplex of M and $R_{i+1} \setminus R_i$ is a PL-subcomplex of $M \setminus R_i$, $i = 1, \dots, n-1$. Let us show that the identity map of Y can be arbitrarily closely approximated by a homeomorphism $g : Y \rightarrow Y$ such that $g(y) = y$ for every $y \in Y \setminus M$, $g(R)$ is a closed rationally presented subset of M and g restricted to $M \setminus R$ is a PL-homeomorphism to $M \setminus g(R)$ sending $\mathcal{P}$ to a rational decomposition of $M \setminus g(R)$ (note that the rational structure of M induces the corresponding rational structure on open subsets of M such that the inclusions are rational maps).

Approximate the identity map of Y by a homeomorphism $g_1 : Y \rightarrow Y$ such that g_1 does not move the points of $Y \setminus M$, g_1 restricted to M is a PL-homeomorphism and $g_1(R_1)$ is a rational subcomplex of M . Approximate g_1 by a map $g_2 : Y \rightarrow Y$ such that g_2 does not move the points of $(Y \setminus M) \cup g_1(R_1)$, g_2 restricted to $M \setminus g_1(R_1)$ is a PL-homeomorphism and $g_2(g_1(R_2 \setminus R_1))$ is a rational subcomplex of $M \setminus g_1(R_1)$. Proceed by induction and construct for every $i = 1, 2, \dots, n-1$ an approximation of the identity map of Y by a homeomorphism $g_{i+1} : Y \rightarrow Y$ such that g_{i+1} does not move the points of $(Y \setminus M) \cup (g_i \circ \dots \circ g_1)(R_i)$, g_{i+1} restricted to $M \setminus (g_i \circ \dots \circ g_1)(R_i)$ is a PL-homeomorphism and $(g_{i+1} \circ \dots \circ g_1)(R_{i+1} \setminus R_i)$ is a rational subcomplex of $M \setminus (g_i \circ \dots \circ g_1)(R_i)$. And finally construct g_n with the additional property that $g = g_n \circ \dots \circ g_1$ sends $\mathcal{P}$ to a rational decomposition of $M \setminus g(R)$. Then g will have the required properties.

3 Proof of Theorem 1.2

Let A be a closed subset of a space X . A collection $\mathcal{C}$ of subsets $X \setminus A$ is said to properly approach A if for every sequence C_j of elements of $\mathcal{C}$ such that there is a sequence of points $x_j \in C_j$ converging to a point $a \in A$ we have that $\lim_{j \rightarrow \infty} C_j = a$ (that is $\lim_{j \rightarrow \infty} y_j = a$ for every sequence $\{y_j\}$ with $y_j \in C_j$). Note that if $\mathcal{C}$ properly approaches A then the closure in X of each $C \in \mathcal{C}$ does not intersect A . Also note that if collections $\mathcal{C}$ and $\mathcal{C}'$ of subsets of $X \setminus A$ properly approach A then $\text{st}(\mathcal{C}, \mathcal{C}')$ properly approaches A as well.

Proposition 3.1 *Let X be an n -dimensional Nöbeling space, let A be a Z -set in X and let $\mathcal{C}$ be a cover of $X \setminus A$ that properly approaches A . Then for every $C \in \mathcal{C}$ there is an open set $C \subset V_C \subset X \setminus A$ such that the inclusion $C \subset V_C$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$ and $\mathcal{V} = \{V_C : C \in \mathcal{C}\}$ properly approaches A .*

Proof. Fix a metric d on X such that $d(x, y) \leq 1$ for every $x, y \in X$. If $C \in \mathcal{C}$ is not a singleton denote $d_C = \inf$ of $\text{diam} G$ for open subsets G of X such that $C \subset G$ and the inclusion $C \subset G$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$ and if $C \in \mathcal{C}$ is a singleton define $d_C = d(C, A)$. For every $C \in \mathcal{C}$ fix an open set G_C such that $\text{diam} G_C \leq 2d_C$ and the inclusion $C \subset G_C$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$.

Let $\mathcal{U}$ be an open cover of $X \setminus A$ which properly approaches A . Denote $\mathcal{C}_n = \{C \in \mathcal{C} : 1/n + 1 < d_C \leq 1/n\}$. Then the closure of the union of the elements of $\mathcal{C}_n$ does not intersect A . For a given n approximate the identity map of X by a closed embedding $e_n : X \rightarrow X$ such that $e_n(X) \subset X \setminus A$ and for $C \in \mathcal{C}_n$, $\text{diam}(e_n(G_C)) \leq \text{diam} G_C + 1/n$, $e_n(C) \subset \text{st}(C, \mathcal{U})$ and every map $f : S^p \rightarrow C$ from a p -dimensional sphere S^p , $p \leq n - 1$ can be homotoped into $e_n(C)$ inside $\text{st}(C, \mathcal{U})$. Denote $Y_C = \text{st}(C, \mathcal{U}) \cup e_n(G_C)$ for $C \in \mathcal{C}_n$.

Then the inclusion $C \subset Y_C$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$ and the cover $\{Y_C : C \in \mathcal{C}\}$ of $X \setminus A$ properly approaches A . Enlarge the sets Y_C to open sets V_C such that $\mathcal{V} = \{V_C : C \in \mathcal{C}\}$ properly approaches A . Then $\mathcal{V}$ has the required properties. $\square$

Suppose that A is a closed subset of X and $X \setminus A$ is embedded as a dense subset in M . Let us show that there is an open subset V of M containing $X \setminus A$ for which there is a space Y such that X and V embed into Y such that $Y = X \cup V$, $X \setminus A = X \cap V$, $A = Y \setminus V$ and A is closed in Y .

Embed X into a Hilbert space and let a metric $d(x, y) = |x - y|$ on X comes from a norm in H . For every $x \in X \setminus A$ define $W(x) = \{y \in X : d(x, y) < d(x, A)/2\}$ and let $\mathcal{W} = \{W(x) : x \in X \setminus A\}$. Refine $\mathcal{W}$ by a locally finite open cover $\mathcal{G}$ of $X \setminus A$. For every $G \in \mathcal{G}$ pick up a point $x_G \in X \setminus A$ such that $G \subset W(x_G)$ and observe that if x belongs to $G \in \mathcal{G}$ then $d(x_G, A)/2 \leq d(x, A) \leq 3d(x_G, A)/2$ and $d(x_G, x) \leq d(x, A)$. Extend each $G \in \mathcal{G}$ to an open subset V_G of M such that $G = (X \setminus A) \cap V_G$ and $\mathcal{V} = \{V_G : G \in \mathcal{G}\}$ is locally finite in $V = \bigcup \{V_G : G \in \mathcal{G}\}$. Let $\{f_G : G \in \mathcal{G}\}$ be a partition of unity subordinated to $\mathcal{V}$. Set $p_G = (x_G, d(x_G, A)) \in H \times (0, \infty)$ and define a map $\phi_V : V \rightarrow H \times (0, \infty)$ by

$\phi_V(x) = \sum_{G \in \mathcal{G}} f_G(x) p_G$. Identify H with $H \times 0 \subset H \times [0, \infty)$ and let $\mathcal{U}$ be an open cover of $H \times (0, \infty)$ that properly approaches H . Approximate ϕ_V by a $\mathcal{U}$ -close embedding $h_V : V \rightarrow H \times (0, \infty)$. Define a set Y as the free union of A and V . Then X is a subset of Y . Denote by h_A the inclusion of A into H and define a function $h : Y \rightarrow H \times [0, \infty)$ such that $h|_A = h_A$ and $h|_V = h_V$.

Let us show that $h|_X$ is an embedding of X into $H \times [0, \infty)$. To show this we need to verify that for a sequence x_j in $X \setminus A$ and $a \in A$, $\lim x_j = a$ if and only if $\lim h_V(x_j) = a$. Since $\mathcal{U}$ properly approaches H , $\lim h_V(x_j) = a$ is equivalent to $\lim \phi_V(x_j) = a$. For every $G \in \mathcal{G}$ containing x_j we have that $d(x_j, A)/2 \leq d(x_G, A) \leq 3d(x_j, A)/2$ and $d(x_G, x_j) \leq d(x_j, A)$. Then $d(x_j, A)/2 \leq \sum_G f_G(x_j) d(x_G, A) \leq 3d(x_j, A)/2$ and $|\sum_G f_G(x_j) x_G - x_j| \leq d(x_j, A)$. Assume that $\lim x_j = a$. Then $\lim d(x_j, A) = 0$ and hence $\lim_j \sum_G f_G(x_j) d(x_G, A) = 0$ and $\lim_j \sum_G f_G(x_j) x_G = \lim x_j = a$. Thus $\lim \phi_V(x_j) = a$. Now assume $\lim \phi_V(x_j) = a$. Then $\lim_j \sum_G f_G(x_j) d(x_G, A) = 0$ and $\lim_j \sum_G f_G(x_j) x_G = a$. This implies $\lim d(x_j, A) = 0$ and $\lim_j \sum_G f_G(x_j) x_G = \lim x_j = a$. Thus we showed that h embeds X into $H \times [0, \infty)$. Now we can transfer the topology of $h(Y)$ to Y and get the required space.

Let M_i be an l_i -co-connected triangulated manifold, $i = 1, 2$ such that $m_i \geq 2(m_i - l_i) + 3$ and $m_1 - l_1 = m_2 - l_2$ for $m_i = \dim M_i$. Denote $k_i = l_i - 2$ and $n = m_i - l_i + 1$. Then by Theorem 1.1, $M_i(k_i)$ is an n -dimensional Nöbeling space. Assume that X_i and A_i satisfy the assumptions of Theorem 1.2, let $f : A_1 \rightarrow A_2$ be a homeomorphism and let $X_i \setminus A_i$ be homeomorphic to $M_i(k_i)$. Identify $X_i \setminus A_i$ with $M_i(k_i)$ and, by the property proved above, replace M_i by an open subset of M_i containing $M_i(k_i)$ and assume that X_i and M_i are subspaces of a space Y_i such that $Y_i = X_i \cup M_i$, $X_i \setminus A_i = X_i \cap M_i$, $A_i = Y_i \setminus M_i$ and A_i is closed in Y_i .

Let $\mu : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ be a one-to-one correspondence of between collections of subsets of M_1 and M_2 respectively. We say that μ agrees with $f : A_1 \rightarrow A_2$ if for every sequence $C_j, j = 1, 2, \dots$ of elements of $\mathcal{C}_1$, C_j converge to $a \in A_1$ in Y_1 (as $j \rightarrow \infty$) if and only if $\mu(C_j)$ converge to $f(a)$ in Y_2 . It is easy to check that if $\mathcal{C}_1$ and $\mathcal{C}_2$ properly approach A_1 and A_2 respectively, μ agrees with f and a collection $\mathcal{C}$ of subsets of M_1 properly approaches A_1 then $\mu(\text{st}(\mathcal{C}, \mathcal{C}_1))$ properly approaches A_2 where $\mu(\text{st}(\mathcal{C}, \mathcal{C}_1)) = \{\mu(\text{st}(C, \mathcal{C}_1)) : C \in \mathcal{C}\}$ and $\mu(\text{st}(C, \mathcal{C}_1)) = \cup\{\mu(C') : C' \in \mathcal{C}_1, C' \cap C \neq \emptyset\}$.

Let M be a triangulated space. It follows from the interpretation of $M(k)$ given in Introduction that if P is either a rational subcomplex or an open subset of a triangulated space M then $P \cap M(k) = P(k)$ where P is considered with the rational structure for which the inclusion of P into M is a rational map (as a matter of fact this is true for every rational subspace P of M however we will not use this fact).

The poof of Theorem 1.2 is based on

Proposition 3.2 *Let $M_i, A_i, X_i, Y_i, k_i, l_i, m_i$ and f be as above and let $\mathcal{P}_1$ be an l_1 -co-connected rational partition of an open submanifold M'_1 of M_1 such that $M_1(k_1) \subset M'_1$, $\mathcal{P}_1$ has no non-empty intersections of $\dim \leq l_1 - 2$ and $\mathcal{P}_1$ properly approaches A_1 . Then there are an open submanifold M'_2 of M_2 and an l_2 -co-connected rational partition of M'_2 such*

that $M_2(k_2) \subset M'_2$, $\mathcal{P}_2$ properly approaches A_2 and $\mathcal{P}_2$ admits a matching $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ which agrees with $f : A_1 \longrightarrow A_2$.

Let us show how Theorem 1.2 can be derived from Proposition 3.2.

Proof of Theorem 1.2. Assume that theorem holds in $\dim \leq n - 1$ and let us prove it in $\dim = n$. Fix complete metrics on X_1 and X_2 and let $\epsilon > 0$. Take a locally finite cover $\mathcal{C}$ of $X_1 \setminus A_1$ of mesh $\mathcal{C} < \epsilon$ such that $\mathcal{C}$ properly approaches A_1 . Then one can find an open subset M'_1 of M_1 such that $M_1(k_1) \subset M'_1$ and the closures in M'_1 of the elements of $\mathcal{C}$ form a locally finite cover of M'_1 . Now there is a rational triangulation of M'_1 such that the decomposition of M'_1 formed from the m_1 -dimensional simplexes of the triangulation properly approaches A_1 and the diameter of the intersection of every m_1 -dimensional simplex with $M_1(k_1)$ is $< 2\epsilon$ with respect to the complete metric fixed in X_1 . Define $\mathcal{P}_1$ as the stars of the vertices of this triangulation with respect to its first barycentric subdivision. Then $\mathcal{P}_1$ is a rational partition of M'_1 which properly approaches A_1 . Note that the intersections of $\mathcal{P}_1$ of $\dim \leq l_1 - 2$ do not intersect $M_1(k_1)$ and therefore by removing them from M'_1 and from the elements of $\mathcal{P}_1$ we may assume that $\mathcal{P}_1$ has no non-empty intersections of $\dim \leq l_1 - 2$. Note all the intersections of $\mathcal{P}_1$ remain contractible and hence $\mathcal{P}_1$ is l_1 -co-connected.

By Proposition 3.2 there are an open submanifold M'_2 of M_2 and an l_2 -co-connected rational partition of M'_2 such that $M_2(k_2) \subset M'_2$, $\mathcal{P}_2$ properly approaches A_2 and $\mathcal{P}_2$ admits a matching $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ which agrees with $f : A_1 \longrightarrow A_2$.

We are going to construct homeomorphisms $f_P : P(k_1) \longrightarrow \mu(P)(k_2)$ for all finite intersections P of $\mathcal{P}_1$ of $\dim \leq m_1 - 1$ which will agree on the common intersections. Let $t \leq m_1 - 1$ and assume that for every finite intersection P of $\dim \leq t - 1$ we already constructed f_P . Take an intersection P of $\mathcal{P}_1$ such that $\dim P = t$. By Theorem 1.1, $P(k_1)$ is a Nöbeling space of $\dim \leq n - 1$. Note for the union P' of the intersections of $\mathcal{P}_1$ of $\dim < t$ that are contained in P , P' lies in ∂P and hence $P'(k_1)$ is a Z -set in $P(k_1)$. Define the homeomorphism $f_{P'} : P'(k_1) \longrightarrow \mu(P')(k_2)$ by the homeomorphisms of the intersections forming P' . According to our assumption Theorem 1.2 holds in $\dim \leq n - 1$. Therefore $f_{P'}$ can be extended to a homeomorphism $f_P : P(k_1) \longrightarrow \mu(P)(k_2)$.

Recall that $\mathcal{P}_1$ and $\mathcal{P}_2$ properly approach A_1 and A_2 respectively and the matching μ agrees with f . Then a homeomorphism extending f between X_1 and X_2 can be obtain by pasting homeomorphisms from $P(k_1)$ to $\mu(P)(k_2)$ for $P \in \mathcal{P}_1$ which extend already defined homeomorphisms on intersections of $\mathcal{P}_1$ of $\dim \leq m_1 - 1$.

Fix $P \in \mathcal{P}_1$ and let P' be the union of the intersections of $\mathcal{P}_1$ of $\dim \leq m_1 - 1$ that are contained in P . The homeomorphism $f_{P'} : P'(k_1) \longrightarrow \mu(P')(k_2)$ is a homeomorphism of Z -subsets of $P(k_1)$ and $\mu(P)(k_2)$ respectively. Therefore we can repeat for P and $\mu(P)$ the same procedure that we did for M_1 and M_2 but this time in the opposite direction from $\mu(P)$ to P first "splitting" $\mu(P)$ into small pieces and then defining the corresponding splitting of P .

Thus going back and force we are able after each iteration to extend f to a partial homeomorphism of bigger and bigger parts of X_1 and X_2 and simultaneously to restrict for each point of X_1 and X_2 the set to which this point can be sent under a possible extension of f . Finally, passing to the limit and using the completeness of X_1 and X_2 we

get the desired homeomorphism. $\square$

Proof of Proposition 3.2. The proof of the proposition splits into two independent parts.

Constructing an initial partition $\mathcal{P}_2$. Let $P_0, \dots, P_t$ be distinct elements of $\mathcal{P}_1$ and let $P = P_0 \cap \dots \cap P_t$. Denote by Δ_P a t -dimensional simplex with vertices $v_0, \dots, v_t$ and for a simplex $\Delta' \subset \Delta_P$ denote by $V_{\Delta'}$ the set of vertices of Δ' . A map $e_P : \Delta_P \rightarrow M'_1$ is called a map witnessing the intersection P if $e_P(\Delta_P) \subset \text{Int}(P_0 \cup \dots \cup P_t)$, $e_P(v_i) \in \text{Int}P_i$ and P_i does not intersect $e(\Delta_P^i)$ where Δ_P^i is the face of Δ_P spanned by the vertices $\{v_0, \dots, v_t\} \setminus \{v_i\}$, $i = 0, \dots, t$. Note that if there is a map e_P witnessing the intersection P then $P \neq \emptyset$. Indeed, aiming at a contradiction assume that $P = \emptyset$. Enlarge each $e^{-1}(P_i)$ to an open subset $e^{-1}(P_i) \subset G_i$ of Δ_P such that G_i does not meet Δ_P^i and $G_0 \cap \dots \cap G_t = \emptyset$. Let $f_0, f_1, \dots, f_t$ be a partition of unity subordinated to the cover $\mathcal{G} = \{G_0, \dots, G_t\}$ of Δ_P . Define $f : \Delta_P \rightarrow \Delta_P$ by $f(x) = f_0(x)v_0 + \dots + f_t(x)v_t$, $x \in \Delta_P$. Then $f(\Delta_P) \subset \partial\Delta_P$ and, for every i and $x \in \Delta_P^i$, $f(x) \in \Delta_P^i$. Hence f restricted to $\partial\Delta_P$ is homotopic to the identity map of $\partial\Delta_P$ and this contradicts the fact that $\partial\Delta_P$ is not a retract of Δ_P .

Assume that P is a non-empty intersection and let us show how to construct a map witnessing the intersection P . Let $F_i \subset \Delta_P$ be the star of v_i with respect to the first barycentric subdivision of Δ_P . For a simplex $\Delta' \subset \Delta_P$ denote $C_{\Delta'} = \cap \{F_i : v_i \in V_{\Delta'}\} \subset \Delta_P$, $P_{\Delta'} = \cap \{P_i : v_i \in V_{\Delta'}\} \subset M'_1$ and let $C_i = \cup \{C_{\Delta'} : \Delta' \subset \Delta_P, \dim \Delta' \geq i\}$. Note that $C_{\Delta'}$ is a singleton and $P_{\Delta'} = P$ for $\Delta' = \Delta_P$. Fix a point $x_P \in \text{Int}P$ and let W be an open neighborhood of x in M'_1 such that $P_{\Delta'} \cap W$ is contractible for every $\Delta' \subset \Delta_P$. We will construct an embedding $e_P : \Delta_P \rightarrow M'_1$ such that $e_P(C_{\Delta'}) \subset P_{\Delta'} \cap W$ for every $\Delta' \subset \Delta_P$ and $e(C_{\Delta'} \setminus C_i) \subset \text{Int}(P_{\Delta'} \cap W)$ for every i and $\Delta' \subset \Delta_P$ such that $\dim \Delta' = i - 1$. The map e_P is constructed by induction on $i = t, t - 1, \dots$ as follows.

Set $e_P(C_{\Delta_P}) = x_P \in \text{Int}(P \cap W)$ and assume that e_P is already defined on C_i such that the required properties are satisfied. Then for an $(i - 1)$ -dimensional simplex $\Delta' \subset \Delta_P$, $e_P(C_i \cap C_{\Delta'}) \subset \partial(P_{\Delta'} \cap W)$ and, since $P_{\Delta'} \cap W$ is contractible, e_P can be extended over $C_{\Delta'}$ such that $e_P(C_{\Delta'}) \subset P_{\Delta'} \cap W$ and $e_P(C_{\Delta'} \setminus C_i) \subset \text{Int}(P_{\Delta'} \cap W)$. This way we extend e_P over C_{i-1} and the construction of e_P is completed. It is easy to see that e_P witnesses the intersection P . Since $\mathcal{P}_1$ has no finite intersection of $\dim \leq l_1 - 2$, $\dim \Delta_P = t \leq m_1 - l_1 + 1 = n$. Then replacing e_P by an approximation which also witnesses the intersection P we may assume that $e_P(\Delta_P) \subset M_1(k_1)$. Note that taking W sufficiently close to x_P we can construct e_P such that the images of $e_P(\Delta_P)$ for all finite intersections of $\mathcal{P}_1$ will form a discrete family in M'_1 .

Fix a collection of maps e_P with images contained in $M_1(k_1)$ and forming a discrete family in M'_1 that witnesses all the intersections of $\mathcal{P}_1$. Extend $f : A_1 \rightarrow A_2$ to a closed embedding $g_1 : X_1 \rightarrow X_2$ such that $g_1(X_1)$ is a Z -set in X_2 . Approximate the map $g_1^{-1}|_{\dots} : g_1(X_1) \rightarrow X_1$ by a closed embedding $g_2 : g_1(X_1) \rightarrow X_1$ such that g_2 restricted to A_2 coincides with f^{-1} , $g_2(g_1(X_1))$ is a Z -set in X_1 and for every finite intersection P of $\mathcal{P}_1$ the map $g_2 \circ g_1 \circ e_P : \Delta_P \rightarrow X_1 \setminus A_1 = M_1(k_1) \subset M'_1$ also witnesses the intersection

P . Finally extend g_2 to a closed embedding $g : X_2 \longrightarrow X_1$.

Then one can find an open subset M'_2 of M_2 such that $M_2(k_2) \subset M'_2$ and M'_2 admits a rational triangulation $\mathcal{T}$ such that 1) $\mathcal{T}$ properly approaches A_2 , 2) for every intersection $P = P_0 \cap \dots \cap P_t$ of distinct elements $P_0, \dots, P_t$ of $\mathcal{P}_1$ and the map $g_1 \circ e_P : \Delta_P = [v_0, \dots, v_t] \longrightarrow M_2(k_2) \subset M'_2$ every simplex of $\mathcal{T}$ intersecting $g_1(e_P(\Delta_P))$ does not intersect $g^{-1}((M'_1 \setminus \text{Int}(P_0 \cup \dots \cup P_t))(k_1))$, every simplex of $\mathcal{T}$ intersecting $g^{-1}(P_i(k_1))$ does not intersect $g_1(e_P(\Delta_P^i))$ and every simplex of $\mathcal{T}$ containing $g_1(e_P(v_i))$ is contained in $g^{-1}((\text{Int} P_i)(k_1))$ and 3) for every finite intersections P and P' of $\mathcal{P}_1$ such that $P \cap P' = \emptyset$, every simplex of $\mathcal{T}$ intersecting $g^{-1}(P(k_1))$ does not intersect $g^{-1}(P'(k_1))$.

Let $\mathcal{B}$ be the partition of M'_2 formed by the stars of vertices of $\mathcal{T}$ with respect to the first barycentric subdivision of $\mathcal{T}$. Arrange the elements of $\mathcal{P}_1 = \{P'_1, P'_2, \dots\}$ into a sequence and define $\mu(P'_1) =$ the union of the elements of $\mathcal{B}$ that intersect $g^{-1}(P'_1(k_1))$, $\mu(P'_2) =$ the union of the elements of $\mathcal{B}$ that intersect $g^{-1}(P'_2(k_1))$ but do not intersect $g^{-1}(P'_1(k_1))$, $\mu(P'_3) =$ the union of the elements of $\mathcal{B}$ that intersect $g^{-1}(P'_3(k_1))$ but do not intersect $g^{-1}(P'_1(k_1)) \cup g^{-1}(P'_2(k_1))$ and so on. Denote $\mathcal{P}_2 = \{\mu(P'_1), \mu(P'_2), \dots\}$. By 2.2, $\mathcal{P}_2$ is a partition of M'_2 .

Then the property 2) guarantees that for every $P \in \mathcal{P}_1$, $g_1(e_P(\Delta_P)) \subset \mu(P)$ and therefore $\mu(P) \neq \emptyset$ and for every non-empty intersection $P = P_0 \cap \dots \cap P_t$ of distinct elements $P_0, \dots, P_t \in \mathcal{P}_1$, the map $g_1 \circ e_P$ witnesses the intersection of $\mu(P_0), \dots, \mu(P_t)$ and therefore $\mu(P_0) \cap \dots \cap \mu(P_t) \neq \emptyset$. The property 3) implies that we do not create additional intersections in $\mathcal{P}_2$ and hence $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ is a matching of partitions on M'_1 and M'_2 respectively. From the property 1) we conclude that $\mathcal{P}_2$ properly approaches A_2 and μ agrees with f .

Improving connectivity of $\mathcal{P}_2$. Without loss of generality we may replace M_1 and M_2 by M'_1 and M'_2 respectively and assume that $M_1 = M'_1$ and $M_2 = M'_2$ for the partition $\mathcal{P}_1$ and the initial partition $\mathcal{P}_2$. Let us show that using 2.10 we can modify $\mathcal{P}_2$ into the required l_2 -co-connected rational partition. The construction of 2.10 is a combination of 2.8 and 2.9, and 2.8 consists of 2.7 applied finitely many times. Abusing the notation we always denote by M_2 , F and $\mathcal{P}_2$ the manifold, the PL-subcomplex of M_2 and the decomposition of M_2 which are the input of 2.7 and 2.9 (=the output the previous applications of 2.7 and 2.9) and by M'_2 , F' and $\mathcal{P}'_2$ the output of 2.7 and 2.9 (=the modifications of M_2 , F and $\mathcal{P}_2$ that are obtained after applying 2.7 and 2.9), and we denote again by μ the correspondence $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}'_2$ which is the composition of $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ with the natural one-to-one correspondence between $\mathcal{P}_2$ and its modification $\mathcal{P}'_2$ (defined by sending each element of $\mathcal{P}_2$ to its modification). Thus we start with the initial partition partition $\mathcal{P}_2$ that during 2.10 turns into a decomposition forming a partition on $U = M_2 \setminus F$ and, gradually reducing the dimension of F , $\mathcal{P}_2$ returns to be a partition at the end of 2.10. Recall that after applying 2.7 and 2.9 we always obtain an open subset M'_2 of M_2 such that $R = M_2 \setminus M'_2$ is a PL-presented subset of M_2 with $\dim R \leq n = m_2 - l_2 + 1 \leq k_2$. By 2.11 we can choose a homeomorphism of $h : A_2 \cup M_2 \longrightarrow A_2 \cup M_2$ such that h is arbitrarily close to the identity map, $h(a) = a$ for $a \in A_2$, $h(R)$ is a rationally presented closed subset of M_2 , $h(F')$ is a rational subcomplex of $h(M'_2)$, $h(\mathcal{P}'_2)$ is a rational decomposition of $h(M'_2)$

and h restricted to $M_2 \setminus R$ is a PL-homeomorphism to $M_2 \setminus h(R)$. Then replacing M'_2 , F' and $\mathcal{P}'_2$ by $h(M'_2)$, $h(F')$ and $h(\mathcal{P}'_2)$ respectively we can always assume that $M_2(k_2) \subset M'_2$, F' is a rational subcomplex of M'_2 and $\mathcal{P}'_2$ is a rational decomposition. Thus at the end of 2.10 we get that $\dim F' \leq n = m_2 - l_2 + 1 \leq k_2$ and hence $M_2(k_1) \subset M'_2 \setminus F'$. Now we can replace the final modification M'_2 of M_2 by $M'_2 \setminus F$ and get the required partition on M'_2 .

It is easy to see that 2.7, 2.9 and 2.11 can be applied such that each element of $\mathcal{P}_2$ will intersect its modification. This implies that $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}'_2$ will automatically agree with f provided $\mathcal{P}'_2$ properly approaches A_2 . Thus in order to get the partition required in Proposition 3.2 the only thing we need to take care of is to perform 2.7 and 2.9 preserving the property that the modification of $\mathcal{P}_2$ properly approach A_2 .

Let us first analyze 2.7. We want to carry out 2.7 in such a way that the modification of $\mathcal{P}_2$ (the output of 2.7) will properly approach A_2 . Assume that $\mathcal{P}_2$ properly approaches A_2 . Enlarge $\text{st}(\mathcal{P}_2, \mathcal{P}_2)$ to an open cover $\mathcal{C}$ of M_2 such that $\mathcal{C}$ properly approaches A_2 and $\text{st}(\mathcal{P}_2, \mathcal{P}_2)$ refines $\mathcal{C}$. By Proposition 3.1 there is an open cover $\mathcal{V}$ of $M_2(k_2)$ such that $\mathcal{V}$ properly approaches A_2 , $\mathcal{C}|_{M_2(k_2)}$ refines $\mathcal{V}$ and for every $C \in \mathcal{C}$ there is $V_C \in \mathcal{V}$ such that $C \cap M_2(k_2) \subset V_C$ and the inclusion $C \cap M_2(k_2) \subset V_C$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$. Then $\mathcal{W} = \text{st}(\mathcal{V}, \mathcal{C})$ is an open cover of M_2 which properly approaches A_2 . Since every map from a space of $\dim \leq n$ to an open subset G of M_2 can be homotoped inside G to a map into $G \cap M_2(k_2)$ we conclude that the inclusion $C \subset \text{st}(V_C, \mathcal{C}) \in \mathcal{W}$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$ for every $C \in \mathcal{C}$. Recall that $\text{st}(\mathcal{P}_2, \mathcal{P}_2)$ refines $\mathcal{C}$. Then the construction 2.7 can be carried out such that the output of 2.7 (the decomposition obtained after applying 2.7) will refine $\text{st}(\mathcal{W}, \mathcal{W})$ and therefore will properly approach A_2 .

Now we will analyze 2.9. Once again we assume that $\mathcal{P}_2$ properly approaches A_2 and we want to carry out 2.9 in such a way that the modification of $\mathcal{P}_2$ (the output of 2.9) will properly approach A_2 as well. The collection $\text{st}(\mathcal{P}_2, \mathcal{P}_2)$ properly approaches A_2 . Since for $P \in \mathcal{P}_2$, $\text{st}(P, \mathcal{P}_2)$ is a union of elements of $\mathcal{P}_2$ we can naturally define the set $\mu^{-1}(\text{st}(P, \mathcal{P}_2))$ as the union of the corresponding elements of $\mathcal{P}_1$. Then the collection $\mu^{-1}(\text{st}(\mathcal{P}_2, \mathcal{P}_2))$ properly approaches A_1 since μ agrees with f . By proposition 3.1 there is an open cover $\mathcal{V}$ of $M_1(k_1)$ such that $\mathcal{V}$ properly approaches A_1 and for every $P \in \mathcal{P}_2$ there is $V_P \in \mathcal{V}$ such that the inclusion $(\mu^{-1}(\text{st}(P, \mathcal{P}_2)))(k_1) \subset V_P$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$. Denote $H_P = \text{st}(V_P, \mu^{-1}(\text{st}(\mathcal{P}_2, \mathcal{P}_2)))$ and $\mathcal{H} = \{H_P : P \in \mathcal{P}_2\} = \text{st}(\mathcal{V}, \mu^{-1}(\text{st}(\mathcal{P}_2, \mathcal{P}_2)))$. Clearly $V_P \subset H_P$, $\mu^{-1}(\text{st}(P, \mathcal{P}_2)) \subset H_P$ and $\mathcal{H}$ properly approaches A_1 . Since $\mu^{-1}(\text{st}(P, \mathcal{P}_2))$ is a rational submanifold of M_1 of $\dim = m_1$, every map from a space of $\dim \leq n$ into $\mu^{-1}(\text{st}(P, \mathcal{P}_2))$ can be homotoped inside $\mu^{-1}(\text{st}(P, \mathcal{P}_2))$ into $(\mu^{-1}(\text{st}(P, \mathcal{P}_2)))(k_1)$ and therefore the inclusion $\mu^{-1}(\text{st}(P, \mathcal{P}_2)) \subset H_P$ induces the zero-homomorphism of the homotopy groups in $\dim \leq n - 1$. Since each element H_P of $\mathcal{H}$ is a union of elements of $\mathcal{P}_1$ we can define $\mu(H_P)$ and $\mu(\mathcal{H}) = \{\mu(H_P) : P \in \mathcal{P}_2\}$. Note that $\mu(\mathcal{H})$ properly approaches A_2 and $\text{st}(\mathcal{P}_2, \mathcal{P}_2)$ refines $\mu(\mathcal{H})$.

Assume that F is a PL-subcomplex of M_2 such that for $U = M_2 \setminus F$ the decomposition $\mathcal{P}_2$ forms on U an l -co-connected partition, $l \geq l_2$ for which $\mu : \mathcal{P}_1 \longrightarrow \mathcal{P}_2$ becomes a matching when $\mathcal{P}_2$ is restricted to U . Use the notation of 2.9 and let Δ be a simplex in

F that we need to absorb. Take an element P of $\mathcal{P}_2$ such that $\Delta \subset P$. Then by 2.3 the inclusion $\text{st}(P, \mathcal{P}_2) \cap U \subset \mu(H_P) \cap U$ induces the zero-homomorphism of the homotopy groups in $\dim \leq m_2 - l$ and therefore S_Δ can be contracted to a point inside $\mu(H_P) \cap U$ and even inside $\mu(H_P) \cap N$ because $\mu(H_P) \cap N$ is a deformation retract of $\mu(H_P) \cap U$. Extend $\mu(\mathcal{H})$ to an open cover $\mathcal{W}$ of M_2 such that $\mathcal{W}$ properly approaches A_2 and $\mu(\mathcal{H})$ refines $\mathcal{W}$. Then 2.9 can be carried out such that the modification of $\mathcal{P}_2$ will refine $\text{st}(\mathcal{W}, \mathcal{W})$ and therefore it will properly approach A_2 . This completes the proof of the proposition. $\square$

4 Appendix

4.1 Extending partitions in the black hole modification

We adopt the notations and the assumptions of 2.4. Recall that T can be represented as the product $T = B^{q+1} \times S$ where S is identified with $a \times S$ for a point a in ∂B^{q+1} . We consider B^{q+1} and S as PL-embedded in Euclidean spaces. It induces the corresponding PL-embedding of T in the product of Euclidean spaces and we refer to these Euclidean spaces when properties of linearity are used, thus we say that a simplex is linear if it is a simplex (linearly spanned by its vertices) in the corresponding Euclidean space and a map of a linear simplex is linear if it is the linear extension of its values on the vertices.

Denote by $p_S : T \rightarrow S$ and $p_B : T \rightarrow B^{q+1}$ the projections. Let $\mathcal{T}_T$ be a triangulation of T such that the simplexes of $\mathcal{T}_T$ are linear, p_S and p_B restricted to every simplex of $\mathcal{T}_T$ are linear and $\mathcal{T}_T$ underlies S and the simplexes of $\mathcal{T}$ contained in T . Let $\mathcal{T}'_T$ be a subdivision of $\mathcal{T}_T$ such that the simplexes of $\mathcal{T}'_T$ are linear and for every simplex Δ of $\mathcal{T}_T$, $p_S(\Delta)$ is a subcomplex of S with respect to $\mathcal{T}'_T$.

Denote by $\mathcal{T}''_T$ the (first) barycentric subdivision of $\mathcal{T}'_T$ and let $\mathcal{B}_T$ be the partition of T formed by the stars of the vertices of $\mathcal{T}'_T$ with respect to $\mathcal{T}''_T$. Then $\mathcal{B}_S = \mathcal{B}_T|_S$ is a partition of S . For a non-empty finite intersection B of $\mathcal{B}_S$ which is the intersection of distinct elements $B_0, \dots, B_t$ of $\mathcal{B}_S$ let the vertex v_B of $\mathcal{T}''_T$ be the barycenter of the simplex of $\mathcal{T}'_T$ spanned by the vertices contained in $B_0 \cup \dots \cup B_t$.

For a finite intersection B of $\mathcal{B}_S$ denote $K_B = p_S^{-1}(B)$, $N_B = p_S^{-1}(\partial B)$ and $M_B = K_B \cap \partial T$. Note that $K_B = B^{q+1} \times B$, $N_B = B^{q+1} \times \partial B$, $M_B = \partial B^{q+1} \times B$, $\partial K_B = N_B \cup M_B$ and $N_B \cap M_B = \partial B^{q+1} \cap \partial B$.

Let us show that $\mathcal{P}_L$ restricted to $M_B \setminus F$ and $\mathcal{P}_L$ restricted to $\partial M_B \setminus F$ are partitions. Assume that $B \in \mathcal{B}_S$ (that is $\dim B = \dim S$). Note that $\mathcal{P}_L$ restricted to $\text{Int} M_B \setminus F_L$ is a partition. Take a point $x \in \partial M_B$ and let Δ_x be the smallest simplex of $\mathcal{T}_T$ containing x . Then $v_B \in p_S(\Delta_x)$ and let $y \in \Delta_x$ be such that $p_S(y) = v_B$. Let G be a neighborhood of x in ∂M_B such that G is contained in the star of y with respect to $\mathcal{T}_T$. Then $(z, t) \rightarrow z(1-t) + ty$, $z \in G, 0 \leq t < 1$ defines a PL-embedding of $G \times [0, 1)$ into M_B such that $G \times [0, 1)$ is a neighborhood of x in M_B and for every simplex $\Delta \in \mathcal{T}_T$ we have that $\Delta \cap (G \times [0, 1)) = (G \cap \Delta) \times [0, 1)$. Recall that $\mathcal{T}_T$ underlies $\mathcal{P}_L|_{\partial T}$ and $F_L \cap \partial T$, and note that $G \times (0, 1)$ is an open subset of ∂T . Therefore $\mathcal{P}_L$ restricted to $(G \times (0, 1)) \setminus F_L$ is a partition. All these facts together imply by 2.2 that $\mathcal{P}_L$ restricted to $(G \times [0, 1)) \setminus F_L$

and $\mathcal{P}_L$ restricted to $G \setminus F_L$ are partitions and hence $\mathcal{P}_L$ restricted to $M_B \setminus F_L$ and $\mathcal{P}_L$ restricted to $\partial M_B \setminus F_L$ are partitions.

Now assume that for a finite intersection $B' \in \mathcal{B}_S$ we already showed that $\mathcal{P}_L$ restricted to $M_{B'} \setminus F_L$ and $\mathcal{P}_L$ restricted to $\partial M_{B'} \setminus F_L$ are partitions and let B be a finite intersection of $\mathcal{B}_S$ such that $B \subset \partial B'$ and $\dim B + 1 = \dim B'$. Then replacing S by $\partial B'$ and T by $B^{q+1} \times \partial B'$ we can repeat the above reasoning to show that $\mathcal{P}_L$ restricted to $M_B \setminus F_L$ and $\mathcal{P}_L$ restricted to $\partial M_B \setminus F_L$ are partitions. Thus we have shown that for every finite intersection of $\mathcal{B}_S$, $\mathcal{P}_L$ restricted to $M_B \setminus F_L$ and to $\partial M_B \setminus F_L$ are partitions.

Let B be a non-empty finite intersection of $\mathcal{B}_S$. Denote by V the set of vertices v of $\mathcal{T}_T''$ such that $v \neq v_B$, $v \in \partial T$ and $[v, v_B]$ is a simplex of $\mathcal{T}_T''$. For each $v \in V$ choose a point v' lying on the interval $[v, v_B]$ connecting v and v_B as follows. If $v \in S$ then set $v' = v$ and note that in this case $v' \in \partial B$. If $v \notin S$ then choose v' such that if $v' \neq v_B$, $v' \neq v$ and $p_S([v', v]) \subset \text{Int} B$. For a simplex Δ of $\mathcal{T}_T''$ spanned by vertices $v_B, v_1, \dots, v_k$, with $v_1, \dots, v_k \in V$ denote by Δ' the simplex spanned by the vertices $v_B, v'_1, \dots, v'_k$ and denote by W_B the union of all such simplexes Δ' . Then W_B is a PL-ball which is a closed neighborhood of v_B in M_B such that $W_B \cap S = M_B \cap S = B$ and for every $x \in \partial W_B$ we have that the interval $[v_B, x]$ is contained in W_B and $[v_B, x] \cap \partial W_B = x$. Let $\pi_B^W : W_B \setminus v_B \rightarrow \partial W_B$ be the radial projection. Using a PL-homeomorphism represent K_B as $W_B \times [0, 1]$ such that W_B is identified with $W_B \times 0$ and extend π_B^W to the radial projection $\pi_B : K_B \setminus v_B = (W_B \times [0, 1]) \setminus v_B \rightarrow \partial K_B \setminus \text{Int} W_B = (\partial W_B \times [0, 1]) \cup (W_B \times 1)$ (π_B is defined with respect to the linear structure of $W_B \times [0, 1]$ induced by the linear structures of W_B and $[0, 1]$).

Let us show that $\mathcal{P}_L$ restricted $M_B \setminus (F_L \cup \text{Int} W_B)$ is a partition. Take $x \in \partial W_B \setminus S$. Then there are an open neighborhood G of x in $\partial W_B \setminus S$ and $\epsilon > 0$ such that $G \times [0, \epsilon)$ is embedded into $M_B \setminus (S \cup \text{Int} W_B)$ by the map $(z, t) \rightarrow z(1+t) - tv_B, z \in G$ such that for every simplex Δ of $\mathcal{T}_T''$, $\Delta \cap (G \times [0, \epsilon)) = (\Delta \cap G) \times [0, \epsilon)$. Recall that $\mathcal{T}_T''$ underlies $\mathcal{P}_L$ and F_L restricted to ∂T and $\mathcal{P}_L$ restricted to $M_B \setminus F_L$ and to $\partial M_B \setminus F_L$ are partitions. Since $G \times (0, \epsilon)$ is an open subset of M_B , $\mathcal{P}_L$ restricted to $(G \times (0, \epsilon)) \setminus F_L$ is a partition and therefore by 2.2, $\mathcal{P}_L$ restricted to $G \setminus F_L$ is a partition. Then, again by 2.2, $\mathcal{P}_L$ restricted to $(G \times [0, \epsilon)) \setminus F_L$ is a partition and hence $\mathcal{P}_L$ restricted $M_B \setminus (F_L \cup \text{Int} W_B)$ is a partition (recall that $S \subset F_L$).

For every $A \in \mathcal{A}$ and every finite intersection B of $\mathcal{B}_S$ define sets A^B and $A^{\partial B}$ as follows. For $B = \emptyset$ set $A^\emptyset = A$ if $A \cap T = \emptyset$ and $A^\emptyset = A \cup S$ if $A \cap T \neq \emptyset$, and for $\dim B = 0$ set $A^{\partial B} = A^\emptyset$. Now by induction on $\dim B$ define A^B and $A^{\partial B}$ so that $A^B = A^{\partial B} \cup \pi_B^{-1}(A^{\partial B} \cap (\partial K_B \setminus \text{Int} W_B))$ for $\dim B \geq 0$ and $A^{\partial B} = \cup \{A^{B'} : B' \subset \partial B\}$ for $\dim B \geq 1$. Clearly A^B and $A^{\partial B}$ are PL-subcomplexes of M . Note that if $A \cap T = \emptyset$ then $A^B = A^{\partial B} = A$ for every B .

From the construction of W_B and π_B it follows that for every simplex $\Delta \in \mathcal{T}_T'$ intersecting W_B we have $\pi_B^{-1}(\Delta \cap \partial W_B) \setminus S = (\Delta \cap W_B) \setminus S$. Then it is easy to see that $S^{\partial B} = S^B = S$, $(A^{\partial B} \cap M_B) \setminus S = (A^B \cap M_B) \setminus S = (A \cap M_B) \setminus S$ and hence $(A^{B_1} \cap A^{B_2}) \setminus S = A^{B_1 \cap B_2} \setminus S$, $(A_1^B \cap A_2^B) \setminus S = (A_1 \cap A_2)^B \setminus S$ for every $A, A_1, A_2 \in \mathcal{A}$ and finite intersections B, B_1, B_2 of $\mathcal{B}_S$.

Denote $\mathcal{P}^B = \{P^B : P \in \mathcal{P}_L\}$ and $\mathcal{P}^{\partial B} = \{P^{\partial B} : P \in \mathcal{P}_L\}$. Recall that $S \subset F_L$.

Let us show by induction on $\dim B$ that $\mathcal{P}^B$ restricted to $K_B \setminus F_L^B$ and $\mathcal{P}^{\partial B}$ restricted to $N_B \setminus F_L^{\partial B}$ are partitions.

For $\dim B = 0$, $N_B = \emptyset$ and therefore $\mathcal{P}^{\partial B}$ restricted to $N_B \setminus F_L^{\partial B}$ is vacuously a partition. Let B be a finite intersection of $\mathcal{B}_S$ such that $\mathcal{P}^{\partial B}$ restricted to $N_B \setminus F_L^{\partial B}$ is a partition. Denote $R_B = \partial K_B \setminus \text{Int}W_B$. From the construction of π_B it follows that $K_B \setminus v_B$ is PL-homomorphic to $R_B \times [0, 1)$ such that R_B corresponds to $R_B \times 0$, $\mathcal{P}^B$ restricted to $K_B \setminus v_B$ corresponds to the decomposition $(\mathcal{P}^{\partial B}|_{R_B}) \times [0, 1)$ of $R_B \times [0, 1)$ and $(F_L^B \cap K_B) \setminus v_B$ corresponds to $(F_L^{\partial B} \cap R_B) \times [0, 1)$. Note that $F_L^{\partial B} \cap M_B = F_L \cap M_B$ and $\mathcal{P}^{\partial B}$ restricted to $M_B \setminus S$ coincides with $\mathcal{P}_L$ restricted to $M_B \setminus S$ and recall that $\mathcal{P}_L$ restricted to $\partial M_B \setminus F_L$ and $M_B \setminus (F_L \cup \text{Int}W_B)$ are partitions. Then, by 2.2, $\mathcal{P}^{\partial B}$ restricted to $R_B \setminus F_L^{\partial B}$ is a partition and therefore $\mathcal{P}^B$ restricted to $K_B \setminus F_L^B$ is also a partition.

Now assume B is a finite intersection of $\mathcal{B}_S$ such that $\dim B > 0$ and for every finite intersection B' of $\mathcal{B}_S$ such that $B' \subset \partial B$ we have that $\mathcal{P}^{B'}$ restricted to $K_{B'} \setminus F_L^{B'}$ is a partition. Then, by 2.2, $\mathcal{P}^{\partial B}$ is a partition on $N_B \setminus F_L^{\partial B}$ since $\{K_{B'} : B' \text{ is a finite intersection of } \mathcal{B}_S \text{ such that } B' \subset \partial B\}$ is a partition of N_B . The induction is completed.

Thus we have shown that $\mathcal{P}^B$ restricted to $K_B \setminus F_L^B$ is a partition for every finite intersection B of $\mathcal{B}_S$. For $A \in \mathcal{A}$ denote $A^\pi = \cup \{A^B : B \in \mathcal{B}_S\}$ and define $\mathcal{P}_L^\pi = \{P^\pi : P \in \mathcal{P}_L\}$. Note that $F_L^B = K_B \cap F_L$ and $\mathcal{P}_L^\pi$ restricted to K_B coincides with $\mathcal{P}^B$. Then, by 2.2, $\mathcal{P}_L^\pi$ is a partition on $T \setminus F_L^\pi$ since $\{K_B : B \text{ is a finite intersection of } \mathcal{B}_S\}$ is a partition of T . It is obvious that $\mathcal{P}_L^\pi$ restricted to $L \setminus S$ coincides with $\mathcal{P}_L$ restricted to $L \setminus S$ and therefore $\mathcal{P}_L^\pi$ is a decomposition of M which is a partition on $M \setminus F_L^\pi$.

Now let us show that for $A_1, A_2 \in \mathcal{A}$, $(A_1 \setminus A_2) \setminus S$ is a strong deformation retract of $(A_1^\pi \setminus A_2^\pi) \setminus S$. To this end it is enough to show that for every finite intersection B of $\mathcal{B}_S$, $(C'_1 \setminus C'_2) \setminus S$ is a strong deformation retract of $(C_1 \setminus C_2) \setminus S$ where $C'_i = A_i^{\partial B} \cap \partial K_B$ and $C_i = A_i^B \cap K_B$. From the construction it follows that $(C_1 \setminus C_2) \setminus S$ can be topologically represented as $C' \times [0, 1)$ with $C' = (C'_1 \setminus C'_2) \setminus (S \cup \text{Int}W_B)$ such that $(C'_1 \setminus C'_2) \setminus S$ is identified with $(C' \times 0) \cup ((C' \cap \partial W_B) \times [0, 1))$. Note that C' is a space admitting a triangulation for which $C' \cap \partial W_B$ is a subcomplex of C' and therefore $(C' \times 0) \cup ((C' \cap \partial W_B) \times [0, 1))$ is a strong deformation retract of $C' \times [0, 1)$.

Let $\Delta \in \mathcal{T}_T$ be contained in ∂T . Since p_S is linear on Δ and $p_S(\Delta)$ is a subcomplex of $\mathcal{T}'_T$ we can conclude that $\dim \Delta \cap M_B \leq \dim \Delta + \dim B - \dim S$ every finite intersection B of $\mathcal{B}_S$. Then, since $\dim \Delta^B \setminus S \leq \dim(\Delta \cap M_B) \setminus S + 1$ for $\dim B = 0$ and $\dim \Delta^B \setminus S \leq \dim \Delta^{\partial B} \setminus S + 1$ for $\dim B > 0$, we get that $\dim \Delta^\pi \setminus S \leq \dim \Delta \setminus S + 1$. Thus $\dim(A^\pi \cap T) \setminus S \leq \dim(A \cap \partial T) \setminus S + 1$ for every $A \in \mathcal{A}$.

The remaining properties required in 2.4 are easy to verify.

4.2 Digging holes for improving connectivity of an intersection

Let M be a triangulated m -dimensional manifold, F a PL-subcomplex of M and $\mathcal{P}$ a decomposition of M which is a partition on $M \setminus F$. Assume that M is $(q-1)$ -connected, $m \geq 2q+1$ and let $S^{q-1} \subset \text{Int}M \setminus F$ be a PL-embedded q -dimensional sphere so that $\mathcal{P}$ restricted to S^{q-1} is a partition and for every finite intersection P' of $\mathcal{P}$ with $P' \cap S^{q-1} \neq \emptyset$, $S^{q-1} \cap P'$ is properly embedded in $P' \setminus F$ (that is $S^{q-1} \cap \partial(P' \setminus F) = \partial(S^{q-1} \cap P')$). Extend the

embedding of S^q to a PL-embedding of $R^m = R^q \times R^{m-q} \subset \text{Int}M$ such that $S^{q-1} = \partial B^q$ bounds a cube $B^q \subset R^q$ where R^q and R^{m-q} are identified with $R^q \times O \subset R^m$ and $O \times R^{m-q} \subset R^m$.

Fix a triangulation $\mathcal{T}$ of M which is the 2-nd barycentric subdivision of a triangulation that underlies S^{q-1} , F and $\mathcal{P}$. By a simplicial neighborhood of $A \subset M$ in a subcomplex X of M we mean the union of the simplexes of $\mathcal{T}$ intersecting A and contained in X . We may assume that $\mathcal{T}$ is chosen so that the simplicial neighborhood of S^{q-1} in M is contained in R^m . For every finite intersection P' of $\mathcal{P}$ that meets S^{q-1} denote by $G_{P'}$ the simplicial neighborhood of $P' \cap S^{q-1}$ in P' . Note that $G_{P''} = G_{P'} \cap P''$ if $P'' \subset P'$, $G_{P'}$ is a regular neighborhood of $P' \cap S^{q-1}$ in $P' \setminus F$ and $P' \cap S^{q-1}$ is locally flat in $P' \setminus F$ because $\dim P' - \dim P' \cap S^{q-1} = m - (q - 1) \geq q + 2$.

For a block bundle ξ over a cell complex X , written ξ/X , denote by σ_ξ the cells of X , by β_ξ the blocks of the total space $E(\xi)$ and by (σ_ξ, β_ξ) the pairs such that β_ξ is the block over σ_ξ , see [6]. We say that ξ underlies $Y \subset E(\xi)$ if Y is a union of blocks of ξ and ξ underlies a collection $\mathcal{Y}$ of subsets of $E(\xi)$ if ξ underlies each $Y \in \mathcal{Y}$.

For every finite intersection P' of $\mathcal{P}$ that meets S^{q-1} we are going to construct by induction on $\dim P' \cap S^{q-1}$ a block bundle $\xi_{P'}$ over $P' \cap S^{q-1}$ such that $E(\xi_{P'}) = G_{P'}$ and $\xi_{P'}|P'' \cap S^{q-1} = \xi_{P''}$ if $P'' \subset P'$. Obviously such a block bundle exists if $\dim P' \cap S^{q-1} = 0$. Assume that for a finite intersection P' of $\mathcal{P}$ a block bundle $\xi_{P''}$ is already constructed for every finite intersection P'' of $\mathcal{P}$ such that $P'' \subset P'$ and $P'' \neq P'$. Then $\xi_{P''}$ define the corresponding block bundle $\xi_{\partial P'}$ over $\partial(P' \cap S^{q-1})$ and by Theorem 4.3 of [6] this block bundle extends to a block bundle $\xi_{P'}$ over P' such that $E(\xi_{P'}) = G_{P'}$.

Thus we have constructed block bundles $\xi_{P'}$ for every finite intersection P' of $\mathcal{P}$ that meets S^{q-1} and these block bundles define the corresponding block bundle ξ over S^{q-1} .

Let B_1^{m-q} be a cube in R^{m-q} . Fix $\epsilon = 1/2$ and for each cell σ_ξ of S^{q-1} define the pair $(\sigma_\eta, \beta_\eta)$ with $\sigma_\eta = \sigma_\xi$ and $\beta_\eta = \gamma_\xi \times B_1^{m-q}$ where $\gamma_\xi = \{x : x = ts, 1 - \epsilon \leq t \leq 1 + \epsilon, s \in \sigma_\xi\} \subset R^q$. Then the pairs $(\sigma_\eta, \beta_\eta)$ form a block bundle η over S^{q-1} such that $E(\eta)$ underlies a regular neighborhood of S^{q-1} in R^m . Hence by Theorem 4.4 of [6] there is a PL-homeomorphism of R^m realizing an isomorphism of ξ and η . Let $e : R^m \rightarrow M$ be the composition of such PL-homeomorphism with the original embedding of R^m in M . Then replacing the original embedding of R^m into M by the embedding e we may assume that ξ coincides with η .

Fix a triangulation $\mathcal{T}_R$ of R^m such that the simplexes of $\mathcal{T}_R$ are linear in R^m and $\mathcal{T}_R$ underlies $\mathcal{P}|R^m$, $F \cap R^m$ and $B_1^q \times B_1^{m-q}$ where $B_1^q = (1 + \epsilon)B^q$. Let $p : R^m = R^q \times R^{m-q} \rightarrow R^{m-q}$ be the projection and let $a \in \text{Int}B_1^{m-q}$ be such that $a \notin p(\Delta)$ for every $\Delta \in \mathcal{T}_R$ with $\dim p(\Delta) < m - q$ and $a \notin \partial p(\Delta)$ for every $\Delta \in \mathcal{T}_R$ with $\dim p(\Delta) = m - q$. Take a cube B_2^{m-q} such that $a + B_2^{m-q} \subset \text{Int}B_1^{m-q}$ and $a + B_2^{m-q} \subset \text{Int}p(\Delta)$ for every $\Delta \in \mathcal{T}_R$ such that $\Delta \cap (B_1^q \times a) \neq \emptyset$. Then the pairs $(\sigma_\tau, \beta_\tau)$ with $\sigma_\tau = (B_1^q \times a) \cap \Delta$ and $\beta_\tau = (B_1^q \times (a + B_2^{m-q})) \cap \Delta$, $\Delta \in \mathcal{T}_R$ form a block bundle τ over $B_1^q \times a$ and we assume that B_2^{m-q} is so small that $E(\tau) = B_1^q \times (a + B_2^{m-q})$ and $E(\tau|\partial B_1^q) = \partial B_1^q \times (a + B_2^{m-q})$.

By Theorem 1.1 of [6] every block bundle over a cube is trivial. Hence τ is isomorphic to a trivial block bundle δ over B_1^q , that is there is a PL-homeomorphism $h : E(\delta) \rightarrow E(\tau)$ where $E(\delta)$ is the product $E(\delta) = B_1^q \times B_3^{m-q} \subset R^m$ of B_1^q with a cube B_3^{m-q} in R^{m-q}

such that $h(x) = (x, a)$ for every $x \in B_1^q$, $(\sigma_\delta, \beta_\delta)$ is a pair of δ if and only if $(h(\sigma_\delta), h(\beta_\delta))$ is a pair of τ and, for every pair $(\sigma_\delta, \beta_\delta)$ of δ , $\beta_\delta = \sigma_\delta \times B_3^{m-q}$.

Identify $B_1^q \times B_3^{m-q}$ with $h(B_1^q \times B_3^{m-q})$ and let B^{m-q} be a cube lying in $\text{Int}B_3^{m-q}$. Then from the construction it follows that

(*) $P' \cap (B_1^q \times \text{Int}B_3^{m-q}) = (P' \cap B_1^q) \times \text{Int}B_3^{m-q}$ for every $P' \in \mathcal{P}$ and $F \cap (B_1^q \times \text{Int}B_3^{m-q}) = (F \cap B_1^q) \times \text{Int}B_3^{m-q}$.

From the construction it also follows that $\mathcal{P}$ restricted to the sets $\text{Int}B_1^q \setminus \text{Int}B^q$, ∂B^q and $B^q \setminus F$ are partitions. Hence by (*) and 2.2, $\mathcal{P}$ restricted to the sets $(\text{Int}B_1^q \setminus \text{Int}B^q) \times \text{Int}B_3^{m-q}$, $(B^q \setminus F) \times (\text{Int}B_3^{m-q} \setminus \text{Int}B^{m-q})$, $\partial B^q \times (\text{Int}B_3^{m-q} \setminus \text{Int}B^{m-q})$, $\partial B^q \times \partial B^{m-q}$, $\partial B^q \times B^{m-q}$ and $(B^q \setminus F) \times \partial B^{m-q}$ are partitions. Then, once again by 2.2, $\mathcal{P}$ restricted to $\text{Int}(B_1^q \times B_3^{m-q}) \setminus (F \cup (\text{Int}(B^q \times B^{m-q})))$ and $\partial(B^q \times B^{m-q}) \setminus F$ are partitions.

Denote $B^m = B^q \times B^{m-q}$. Thus we get that

(**) $\mathcal{P}$ restricted to $M \setminus (F \cup \text{Int}B^m)$ and $\partial B^m \setminus F$ are partitions.

Let P' be a finite intersection of $\mathcal{P}$. Let us show that

(***) $P' \setminus (F \cup \text{Int}B^m)$ is a deformation retract of $P' \setminus (F \cup \text{Int}B^q)$.

Note the following general fact: for a triangulated space A_1 , a subcomplex A_2 of A_1 and a cube B the space $(A_2 \times B) \cup (A_1 \times \partial B)$ is a strong deformation retract of $(A_2 \times B) \cup (A_1 \times (B \setminus O))$. Then for $A_1 = (P' \cap B^q) \setminus F$ and $A_2 = P' \cap \partial B^q$ we have by (*) that $B^m \cap (P' \setminus (F \cup \text{Int}B^m)) = (A_2 \times B^{m-q}) \cup (A_1 \times \partial B^{m-q})$ and $B^m \cap (P' \setminus (F \cup \text{Int}B^q)) = (A_2 \times B^{m-q}) \cup (A_1 \times (B^{m-q} \setminus O))$. Thus $B^m \cap (P' \setminus (F \cup \text{Int}B^m))$ is a strong deformation retract of $B^m \cap (P' \setminus (F \cup \text{Int}B^q))$ and hence (***) holds.

Now let us compare the embedding of $S^{q-1} = \partial B^q$ into M under h with the embedding of S^{q-1} under e (which coincides with the original embedding of S^{q-1} into M). One can easily observe that there is a PL-homeomorphism $\psi : M \rightarrow M$ such that $\psi \circ h|_{S^{q-1}} = e|_{S^{q-1}}$, ψ does not move the points outside $E(\eta)$, $\psi(P') = P'$ and for every $P' \in \mathcal{P}$. Since ψ does not move the points outside $E(\eta)$ we have that $\psi(F) = F$. Then replacing the embedding of $B_1^q \times B_3^{m-q}$ into M by its composition with ψ we preserve the properties (*), (**) and (***) and obtain that the embedding of S^{q-1} coincides with the original embedding of S^{q-1} .

Thus we can extend the original embedding of S^{q-1} to a PL-embedding of $B^m = B^q \times B^{m-q}$ such that $\mathcal{P}$ restricted to $\partial B^m \setminus F$ and $M \setminus (\text{Int}B^m \cup F)$ are partitions, $S^{q-1} = \partial B^q$, $F \cap B^m = (F \cap B^q) \times B^{m-q}$, $P' \cap B^m = (P' \cap B^q) \times B^{m-q}$ for every $P' \in \mathcal{P}$ and $P' \setminus (F \cup \text{Int}B^m)$ is a deformation retract of $P' \setminus (F \cup \text{Int}B^q)$ for every finite intersection P' of $\mathcal{P}$. It is clear that $B^m \subset R^m \subset \text{Int}M$.

Note that if S^{q-1} can be contracted to a point in an open subset W of M then we may assume that B^m is contained in W .

4.3 Digging holes for absorbing simplexes

We adopt the notations and assumptions of 2.9. Recall that $\mathcal{T}$ is the 2-nd barycentric subdivision of some triangulation $\mathcal{T}_0$ of M underlying $\mathcal{P}$ and consider M as embedded in a Hilbert space by an embedding which is linear on every simplex of $\mathcal{T}_0$.

Fix $\Delta \in \mathcal{T}_F$ and let $S^{q-1} = S_\Delta$ be the link of Δ with respect to $\mathcal{T}$, $S^{m-1} = \partial(\Delta * S_\Delta)$ and b the barycenter of Δ .

Let us show that $\mathcal{P}$ restricted to N , $S^{m-1} \setminus \partial\Delta$ and $(\Delta * S_\Delta) \setminus \Delta$ are partitions. For a point $x \in S^{m-1} \setminus \partial\Delta$ choose an $\epsilon > 0$ and a neighborhood G of x in $S^{m-1} \setminus \partial\Delta$ such that the map $(z, t) \longrightarrow z + tb$, $z \in G$, $t \in (-\epsilon, \epsilon)$ embeds $G \times (-\epsilon, \epsilon)$ into $M \setminus F$ and for every simplex Δ' of $\mathcal{T}_0$ that intersects $G \times (-\epsilon, \epsilon)$ we have that $\Delta' \cap (G \times (-\epsilon, \epsilon)) = (\Delta' \cap G) \times (-\epsilon, \epsilon)$. Then for every $P \in \mathcal{P}$ that intersects $G \times (-\epsilon, \epsilon)$ we have that $P \cap (G \times (-\epsilon, \epsilon)) = (P \cap G) \times (-\epsilon, \epsilon)$. Therefore, by 2.2, $\mathcal{P}$ restricted to G , $G \times (-\epsilon, 0]$ and $G \times [0, \epsilon)$ are partitions and hence $\mathcal{P}$ restricted to $S^{m-1} \setminus \partial\Delta$, $(\Delta * S_\Delta) \setminus \Delta$ and N are partitions as well.

Denote by $\mathcal{T}_\Delta$ the collection of the simplexes of $\mathcal{T}$ intersecting S^{q-1} and containing Δ . Let $\mathcal{T}'$ be the second barycentric subdivision of $\mathcal{T}$. For every $\Delta' \in \mathcal{T}_\Delta$ denote by $G_{\Delta'}$ the simplicial neighborhood of $\Delta' \cap S^{q-1}$ in $\Delta' \cap S^{m-1}$ with respect to $\mathcal{T}'$. In a way similar to constructing $\xi_{P'}$ in 4.2 we construct for every $\Delta' \in \mathcal{T}_\Delta$ a block bundle $\xi_{\Delta'}$ over $\Delta' \cap S^{q-1}$ such that $E(\xi_{\Delta'}) = G_{\Delta'}$ and $\xi_{\Delta'}|(\Delta'' \cap S^{m-1}) = \xi_{\Delta''}$ if $\Delta'' \subset \Delta'$. Then the block bundles $\xi_{\Delta'}$ define the corresponding block bundle ξ over S^{q-1} with $E(\xi)$ being the simplicial neighborhood of S^{q-1} in S^{m-1} with respect to $\mathcal{T}'$. Note that $E(\xi)$ is a regular neighborhood of S^{q-1} in ∂N .

Extend the embedding of S^{q-1} in N to a PL-embedding of a q -dimensional cube B_1^q into N such that $S^{q-1} = \partial B_1^q$ and $B_1^q \cap \partial N = S^{q-1}$ and note that the embedding of B_1^q is locally flat because $m \geq 2q + 1$. Extend the regular neighborhood $E(\xi)$ of S^{q-1} in ∂N to a regular neighborhood of B_1^q in N and by Theorem 4.3 of [6] represent this neighborhood as the total space $E(\nu)$ of a block bundle ν over B_1^q such that $\nu|S^{q-1} = \xi$. Since ν is a trivial block bundle, $E(\nu)$ is PL-homeomorphic to the product $E(\nu) = B_1^q \times B_1^{m-q}$ of B_1^q with an $(m - q)$ -dimensional cube B_1^{m-q} such that for each pair (σ_ν, β_ν) of ν , $\beta_\nu = \sigma_\nu \times B_1^{m-q}$. Fix a triangulation $\mathcal{T}_\nu$ of $E(\nu)$ which underlies $\mathcal{T}$ restricted to $E(\nu)$. In a way similar to constructing the block bundle τ in 4.2 choose a point a in B_1^{m-q} and a cube B_2^{m-q} such that $a + B_2^{m-q} \subset \text{Int} B_1^{m-q}$ and the pairs $(\sigma_\tau, \beta_\tau)$ defined by $\sigma_\tau = \Delta' \cap (B_1^q \times a)$, $\beta_\tau = \Delta' \cap (B_1^q \times (a + B_2^{m-q}))$, $\Delta' \in \mathcal{T}_\nu$ form a block bundle τ over $B_1^q \times a$ with $E(\tau) = B_1^q \times (a + B_2^{m-q})$ and $E(\tau|(\partial(B_1^q \times a))) = E(\tau) \cap (\partial B_1^q \times (a + B_2^{m-q})) (= E(\tau) \cap S^{m-1})$. Then the block bundle τ underlies $\mathcal{P}$ and $\mathcal{T}_\Delta$ restricted to $E(\tau)$.

Note that since the block bundle ν is trivial, the block bundle ξ is trivial as well and $E(\xi) = \partial B_1^q \times B_1^{m-q}$. Then it is obvious that the projection $\partial B_1^q \times a \longrightarrow \partial B_1^q$ extends to a PL-homeomorphism $h : E(\xi) \longrightarrow E(\xi)$ such that $h(x) = x$ for every $x \in \partial E(\xi)$, $h(\beta_\xi) = \beta_\xi$ for every block β_ξ of ξ and there is a PL-isotopy $H : E(\xi) \times [0, 1] \longrightarrow E(\xi) \times [0, 1]$ relative to $\partial E(\xi)$ so that $H_0 = h$. Recall that the barycenter of Δ is denoted by b . Then, since $\partial\Delta \cap E(\xi) = \emptyset$, there is $\epsilon > 0$ such that for every $x \in E(\xi)$, $P \in \mathcal{P}$ and $t \in [0, 1]$ the point $b + (1 + t\epsilon)(x - b)$ belongs to P if and only if x belongs to P . Embed $E(\xi) \times [0, 1]$ into N by sending (x, t) to $b + (1 + t\epsilon)(x - b)$ and define the PL-homeomorphism $g : N \longrightarrow N$ such that g coincides with H on $E(\xi) \times [0, 1]$ and g does not move the points outside $E(\xi) \times [0, 1]$. From the properties described above it is clear that $g(\partial(B_1^q \times a)) = \partial B_1^q$ and the block bundle $g(\tau)$ underlies $\mathcal{P}$ and $\mathcal{T}_\Delta$ restricted to $E(g(\tau))$. Denote $B_2^q = g(B_1^q \times a)$ and $\theta = g(\tau)$.

Thus we get a block bundle θ over a cube B_2^q such that $S^{q-1} = \partial B_2^q = \partial N \cap B_2^q$,

θ underlies $\mathcal{P}$ and $\mathcal{T}_\Delta$ restricted to $E(\theta)$ and $E(\theta|\partial B_2^q) = S^{m-1} \cap E(\theta)$. Then, since θ is trivial, $E(\theta)$ is PL-homomorphic to $B_2^q \times B_3^{m-q}$ such that $\partial N \cap (B_2^q \times B_3^{m-q}) = \partial B_2^q \times B_3^{m-q} \subset S^{m-1}$ and for each pair $(\sigma_\theta, \beta_\theta)$ of θ , $\beta_\theta = \sigma_\theta \times B_3^{m-q}$. Take a cube B_4^{m-q} such that $B_4^{m-q} \subset \text{Int} B_3^{m-q}$ and define the block bundle ρ over B_2^q by restricting the blocks of θ to $E(\rho) = B_2^q \times B_4^{m-q}$. Using the reasoning similar to the one applied for proving the property $(**)$ in 4.2 we can show that $\mathcal{P}$ restricted $N \setminus (B_2^q \times \text{Int} B_4^{m-q})$ and $\partial((\Delta * S_\Delta) \cup (B_2^q \times B_4^{m-q})) \setminus \partial \Delta$ are partitions. It is clear that $E(\rho) \subset \text{Int} M$.

Take a pair $(\sigma_\rho^\Delta, \beta_\rho^\Delta)$ of ρ such that $\dim \beta_\rho^\Delta = m$ and let $P_\Delta \in \mathcal{P}$ be such that $\beta_\rho^\Delta \subset P_\Delta$. Since ρ is a trivial block bundle we can replace ρ by another block bundle subdividing the cells of B_2^q into smaller cells and defining the corresponding blocks using the product structure. This way we may assume that cells of B_2^q are so small that the pair $(\sigma_\rho^\Delta, \beta_\rho^\Delta)$ can be chosen so that $\beta_\rho^\Delta \subset \text{Int} P_\Delta$.

Let the cube $B_3^q \subset \Delta * S_\Delta$ be the join $B_3^q = b * S_\Delta$. Define a block bundle ψ over B_3^q as follows: the pairs $(\sigma_\rho, \beta_\rho)$ of ρ with $\sigma_\rho \subset S^{q-1}$ are pairs of ψ , (b, Δ) is a pair of ψ and the rest of the pairs $(\sigma_\psi, \beta_\psi)$ of ψ are of the form $\sigma_\psi = \Delta' \cap B_3^q$, $\beta_\psi = \Delta'$, $\Delta' \in \mathcal{T}_\Delta$. One can easily check that ψ is indeed a block bundle and $E(\psi) = \Delta * S_\Delta$. Since ψ and ρ over S^{q-1} coincide, they form the corresponding block bundle ϕ over the sphere $B_3^q \cup B_2^q$. Note that $B^q = (B_2^q \cup B_3^q) \setminus \text{Int} \sigma_\rho^\Delta$ is a q -dimensional cube. Define the block bundle η as ψ restricted to B^q . Then (b, Δ) is a pair of η , $b \in \text{Int} B^q$, $\partial B^q = \partial \sigma_\rho^\Delta$, $E(\eta|\partial B^q) \subset \text{Int} P_\Delta$, η underlies $\mathcal{P}$ restricted to $E(\eta)$, $E(\eta) \cap F = \Delta$ and hence η underlies F restricted to $E(\eta)$.

Thus assuming that the center O of B^q is located at b we can represent $E(\eta)$ as the product $B^m = B^q \times B^{m-q}$ such that $\Delta = 0 \times B^{m-q} = F \cap B^m$, $\partial B^q \times B^{m-q} \subset \text{Int} P_\Delta$ and $P \cap B^m = (P \cap B^q) \times B^{m-q}$ for every $P \in \mathcal{P}$ that meets B^m . By the reasoning similar to the one applied to the proof of $(***)$ in 4.2 these properties imply that for every finite intersection P of $\mathcal{P}$, $P \setminus (F \cup \text{Int} B^m)$ is a deformation retract of $P \setminus (F \cup \text{Int} B^q)$. Since $B^m = (\Delta * S_\Delta) \cup ((B_2^q \setminus \text{Int} \sigma_\rho^\Delta) \times B_4^{m-q})$ and $\beta_\rho^\Delta = \sigma_\rho^\Delta \times B_4^{m-q}$ is contained in $\text{Int} P_\Delta$ we derive from the properties of ρ that $\mathcal{P}$ restricted to $M \setminus (F \cup \text{Int} B^m)$ and $\partial B^m \setminus F$ are partitions and $B^m \subset \text{Int} M$.

Note that if S^{q-1} can be contracted to a point in an open subset W of N then the construction can be carried out such that B^m is contained in $(\Delta * S_\Delta) \cup W$.

References

- [1] Ageev, S. Axiomatic method of partitions in theory of Menger and Nöbeling spaces. Preprint (2000), <http://at.yorku.ca/v/a/a/a/87.htm>
- [2] Bestvina, Mladen Characterizing k -dimensional universal Menger compacta. Mem. Amer. Math. Soc. 71 (1988), no. 380, vi+110 pp.
- [3] Chigogidze, A. Inverse spectra. North-Holland Mathematical Library, 53. North-Holland Publishing Co., Amsterdam, 1996. x+421 pp.

- [4] Hudson, J. F. P. Piecewise linear topology. University of Chicago Lecture Notes prepared with the assistance of J. L. Shaneson and J. Lees W. A. Benjamin, Inc., New York-Amsterdam 1969 ix+282 pp
- [5] Kawamura, K.; Levin, M.; Tymchatyn, E. D. A characterization of 1-dimensional Nöbeling spaces. Proceedings of the 12th Summer Conference on General Topology and its Applications (North Bay, ON, 1997). Topology Proc. 22 (1997), Summer, 155–174.
- [6] Rourke, C. P.; Sanderson, B. J. Block bundles. I. Ann. of Math. (2) 87 1968 1–28.
- [7] Rourke, C. P.; Sanderson, B. J. Introduction to piecewise-linear topology. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 69. Springer-Verlag, New York-Heidelberg, 1972. viii+123 pp.

Department of Mathematics
 Ben Gurion University of the Negev
 P.O.B. 653
 Be'er Sheva 84105, ISRAEL
 e-mail: mlevine@math.bgu.ac.il